

**METHODS AND SYSTEMS FOR FACILITATING THE DIAGNOSIS AND
TREATMENT OF SCHIZOPHRENIA**

- [1] This invention was made with United States Government support in the form of Grant Nos. MH45156, MH01489, MH56242, MH53459, and MH45156 from the National Institute of Mental Health. The United States Government may have certain rights in the invention.

FIELD OF THE INVENTION

- [2] The present invention relates generally to the field of neurological and physiological dysfunctions associated with schizophrenia. The invention further relates to the identification, isolation, and cloning of genes which, when mutated or varied, are associated with schizophrenia. The present invention also relates to methods for diagnosing and detecting carriers of the genes and to diagnosis of schizophrenia. The present invention further relates to the construction of animal models of schizophrenia.

BACKGROUND OF THE INVENTION

- [3] Schizophrenia is a serious brain disorder that affects approximately 1% of the human population. The cause of this complex and devastating disease remains elusive, although genetic, nutritional, environmental, and developmental factors have been considered. A combination of clinical, neuroimaging, and postmortem studies have implicated the dorsal prefrontal cortex (PFC) as a prominent site of dysfunction in schizophrenia.

[4] Schizophrenia is typically characterized as a disorder of thinking and cognition, as contrasted to other disorders of mental faculties, such as mood, social behavior, and those affecting learning, memory, and intelligence. Schizophrenia is characterized by psychotic episodes during which an individual may lose the ability to test reality or may have hallucinations, delusions, incoherent thinking, and even disordered memory. There are varying forms of schizophrenia differing in severity, from a schizotypal disorder to a catatonic state. A review of schizophrenia can be found in *Principles of Neural Science*, 3rd ed., 1991, Kandel, Schwartz, and Jessel (Eds.), Connecticut: Appleton & Lange, pp.853-868; of which Chapter 55 is incorporated herein by reference.

[5] Diseases of organ systems, such as those of the heart, lung, and kidney, are usually confirmed by tissue pathology. A demonstrable pathology includes identifying and defining a structural abnormality in the organ, along with an associated alteration in organ function. This type of diagnosis is also utilized in certain neurological diseases. However, there are few psychiatric disorders in which clinical manifestations and symptoms can be correlated with a demonstrable pathology. The majority of mental illnesses are evaluated by observing changes in key behaviors such as thinking, mood, or social behavior. These alterations are difficult to ascertain and nearly impossible to quantify. However, progress is being made in diagnosing mental illness and in determining the neuropathology of mental illnesses.

- [6] The Diagnostic and Statistical Manual of Mental Disorders, Third Edition (DSM-III-R) and the updated DSM-IV, published by the American Psychiatric Association, represent the progress made in providing a basis for objective and rigorous descriptive criteria for categories of psychiatric disorders. While the DSM-III-R is very thorough and detailed, it is also quite lengthy. Thus, the process of reviewing the categories and applying them to data from a patient is also very time-consuming and arduous. In addition, there is no mechanism by which a patient can be diagnosed either as having or being susceptible to schizophrenia prior to the expression of symptoms. Thus, there is a longstanding need for an easy and definitive method for diagnosing schizophrenia. A diagnostic tool that can be applied prior to the expression of symptoms would also have great utility, providing a basis for the development of therapeutic interventions.
- [7] There is strong evidence for a genetic linkage of schizophrenia. Historically, there have been a number of studies on monozygotic twins of schizophrenics that indicated that 30 - 50% of the twins also had schizophrenia. The fact that this number is not 100% indicates that there are other factors involved in this disease process that may protect some of these individuals from the disease. It is apparent from a number of studies that the patterns of inheritance in most forms of schizophrenia are more complex than the classical dominant or recessive Mendelian inheritance. Recently, locus 1q21-22, a chromosome region containing several hundred genes, has been strongly linked to schizophrenia as shown by Brzustowicz et al., *Science*

288, 678-82, 2000, which is hereby incorporated by reference.

[8] Until the 1950's there were no specific, effective treatments for schizophrenia. Antipsychotic drugs were identified in the 1950's, and these drugs were found to produce a dramatic improvement in the psychotic phase of the illness. Reserpine was the first of these drugs to be used and was followed by typical antipsychotic drugs including phenothiazines, the butyrophenones, and the thioxanthenes. A new group of therapeutic drugs, typified by clozapine, has been developed and were referred to as "atypical" antipsychotics. Haloperidol has been employed extensively in the treatment of schizophrenia and is one of the currently preferred options for treatment. When these drugs are taken over the course of at least several weeks, they mitigate or eliminate delusions, hallucinations, and some types of disordered thinking. Maintenance of a patient on these drugs reduces the rate of relapse. Since there is no way of determining if an individual is susceptible to schizophrenia, it is currently unknown if these antipsychotic compounds are useful in the prophylactic treatment of schizophrenia.

[9] Signal transduction is the general process by which cells respond to extracellular signals (e.g. neurotransmitters) through a cascade of biochemical reactions. The first step in this process is the binding of a signaling molecule to a cell membrane receptor that typically leads to the inhibition or activation of an intracellular enzyme. This type of process regulates many cell functions including cell proliferation, differentiation, and gene transcription.

[10] One important mechanism by which signal transduction occurs is through G-proteins. Receptors on the cell surface are coupled intracellularly to a G-protein that becomes activated, when the receptor is occupied by an agonist, by binding to the molecule GTP. Activated G-proteins may influence a large number of cellular processes including voltage-activated calcium channels, adenylate cyclase, and phospholipase C. The G-protein itself is a critical regulator of the pathway by virtue of the fact that GTPase activity in the G-protein eventually hydrolyzes the bound GTP to GDP, restoring the protein to its inactive state. Thus, the G-protein contains a built-in deactivation mechanism for the signaling process.

[11] Recently, an additional regulatory mechanism has been discovered for G-protein signaling that involves a family of mammalian gene products termed regulators of G-protein signaling, or RGS (Druey et al., 1996, Nature 379: 742-746 which is hereby incorporated by reference). RGS molecules play a crucial modulatory role in the G-protein signaling pathway. RGS proteins bind to the GTP-bound $G\alpha$ subunits with a variable $G\alpha$ specificity as a substrate. RGS molecules shorten the GTP binding of the activated $G\alpha$ subunits by acting as GTPase activating proteins (GAPs), accelerating GTP hydrolysis by up to one hundred fold. By the virtue of this GAP action and by making available the GDP-bound $G\alpha$ to re-attach to $\beta\gamma$ dimers, RGS proteins shorten the duration of the intracellular signaling. RGS proteins are expressed in nearly every cell; however, they show a tissue-specific expression across the body and cell type-specific expression in the brain. For example, RGS4 is strongly expressed in the central nervous system, moderately

expressed in the heart, and slightly expressed in skeletal muscle (Nomoto et al., 1997, Biochem. Biophys. Res. Commun. 241(2):281-287 which is herein incorporated by reference).

- [12] Several members of the G-protein signaling pathways, most located downstream of RGS4 modulation, have been implicated in schizophrenia. Gil, Gq and Golf messenger RNA (mRNA) and protein levels all have been reported to be altered in various brain regions of the schizophrenic subjects. Furthermore, changes in expression of adenylate cyclase, phospholipase C, and protein kinases, as well as DARPP (dopamine- and cAMP-regulated phosphoprotein) phosphorylation changes are expected to be influenced by RGS regulation of G α signaling. In addition, RGS modulation changes are expected to have significant effects on the signal transduction effected by neurotransmitters including dopamine, serotonin, GABA, glutamate, and norepinephrine.
- [13] An additional genetic marker of schizophrenia has been identified by Meloni et al. (U.S. patent no. 6,210,879). These investigators found that an allele of the microsatellite HUNTH01 in the tyrosine hydroxylase gene correlated with the expression of schizophrenia. However, the allele only appears to be present in sporadic schizophrenias.
- [14] There has been a long-standing need for a definitive and easy method for diagnosing schizophrenia as well as for an effective treatment with minimal side effects. Further, a need has been recognized in connection with being able to detect schizophrenia prior to the expression of noticeable symptoms.

- [15] A need has been recognized in connection with overcoming the various limitations to the current implementation of a method for diagnosing and assessing the susceptibility to schizophrenia are addressed through the use of the current invention.

SUMMARY OF THE INVENTION

- [16] In accordance with at least one embodiment of the present invention, there is provided a system and method for diagnosing and determining the susceptibility to schizophrenia.
- [17] In summary, one aspect of the present invention provides an isolated and substantially purified DNA sequence corresponding to SEQ ID NOS: 3, 4, 5, 6, 7, 8, and contiguous portions thereof.
- [18] Another aspect of the present invention is a polynucleotide sequence that is complementary to a sequence selected from the group consisting of SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, and contiguous portions thereof.
- [19] A further aspect of the present invention is an expression system comprising a DNA sequence that corresponds to SEQ ID NO:3.
- [20] A yet further aspect of the present invention is a method for diagnosing schizophrenia in a human comprising obtaining a DNA sample comprising a RGS4 gene from a patient and detecting a variation in the RGS4 gene indicating schizophrenia.
- [21] A still further aspect of the present invention is a method for determining the susceptibility to

schizophrenia comprising obtaining from a patient a DNA sample comprising a RGS4 gene and detecting a variation in said RGS4 gene indicating susceptibility to schizophrenia.

[22] An additional aspect of the present invention is a method for diagnosing schizophrenia comprising obtaining from a patient to be tested for schizophrenia a sample of tissue, measuring RGS4 mRNA levels in said sample, and determining if there is a reduced level of RGS4 mRNA in the sample.

[23] A still additional aspect of the present invention is a method of determining susceptibility to schizophrenia comprising obtaining from a patient to be tested for susceptibility to schizophrenia a sample of tissue, measuring RGS4 mRNA levels in said sample, and determining if there is a reduced level of RGS4 mRNA in the sample.

[24] A yet further aspect of the present invention is A method of determining susceptibility to schizophrenia comprising obtaining from a patient to be tested for susceptibility to schizophrenia a sample of tissue, measuring RGS4 protein levels in said sample, and determining if there is a reduced level of RGS4 protein in the sample.

[25] Yet another aspect of the present invention is A method of treating schizophrenia, said method comprising measuring RGS4 protein or mRNA levels in a patient, and altering said RGS4 protein levels to provide the patient with an improved psychiatric function.

[26] Another aspect of the present invention is a kit for diagnosing schizophrenia in a patient, said kit

comprising antibodies to RGS4, and a detector for ascertaining whether said antibodies bind to RGS4 in a sample.

[27] Another aspect of the present invention is a kit for diagnosing schizophrenia in a patient, said kit comprising a detect of RGS4 transcript levels in a patient, and a standard to ascertain altered levels of RGS4 transcript in the patient.

[28] A still further aspect of the present invention is the DNA sequence of SEQ ID NO: 3 containing variations as described in the text below.

[29] A yet further aspect of the present invention is a transgenic mouse whose genome comprises a disruption of the endogenous RGS4 gene, wherein said disruption comprises the insertion of a transgene, and wherein said disruption results in said transgenic mouse not exhibiting normal expression of RGS4 protein.

[30] A still additional aspect of the present invention is a transgenic mouse wherein a transgene comprises a nucleotide sequence that encodes a selectable marker.

[31] These and other embodiments and advantages of the present invention will be better understood with reference to the following figures and detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[32] The present invention and its presently preferred embodiments will be better understood by reference to the detailed disclosure hereinbelow and to the accompanying drawings, wherein:

- [33] **Figure 1A** displays the design of microarray immobilized probes and *in situ* probes for RGS4, wherein numbers on the RGS4 nucleic acid fragments denote nucleotide position in relationship to the RGS4 mRNA, as currently in the NCBI database;
- [34] **Figure 1B** is a pseudocolored intensity view of a single RGS4 feature from the 516 control/547 schizophrenic PFC comparison after a dual-fluorescent hybridization; both images represent the same spot under cy3 and cy5 excitation, respectively; the balanced cy3 signal intensity (c-control subject) was 6.2-fold brighter than the cy5 signal intensity (s-schizophrenic subject);
- [35] **Figure 1C** displays changes in RGS expression in the PFC of schizophrenic and control subjects reported by cDNA microarray analysis;
- [36] **Figure 2A** shows *in situ* hybridization results for PFC RGS4 expression levels which are decreased in 9 of 10 schizophrenic subjects;
- [37] **Figure 2B** shows the *in situ* hybridization data from 10 PFC pairwise comparisons which were quantified using film densitometry;
- [38] **Figure 3A** shows that 632 G-protein signalling-related genes were detected out of 1644 possible detections (274 genes/microarray x six microarrays);
- [39] **Figure 3B** shows that 239 1q21-22 locus-related genes were detected out of 420 possible detections (70 genes/mircoarray x six microarrays); RGS4 contribution to the transcript distribution is denoted by a hatched bar;

- [40] **Figure 4A** shows high power photomicrographs of VC tissue sections from the same matched pair of schizophrenic and matched control subjects represented in FIG. 2A, viewed under darkfield illumination;
- [41] **Figure 4B** shows a graph of 10 supragranular VC SCH pairwise comparisons, in which schizophrenic subjects showed a comparably significant RGS4 transcript reduction to the PFC comparisons;
- [42] **Figure 4C** shows high power photomicrographs of MC tissue sections from the same matched pair of schizophrenic and matched control subjects represented in FIG. 2A, viewed under darkfield illumination;
- [43] **Figure 4D** shows a graph in which schizophrenic subjects across the same 10 subject pairs across the MC had comparably decreased RGS4 expression levels (mean = -34.2%, $F_{1,15} = 10.18$; $p = 0.006$) to VC and PFC;
- [44] **Figure 5** shows a scatter plot of relative RGS4 expression changes across the experimental groups.
- [45] **Figure 6** displays the genomic organization that is derived from available sequences for clone NT_022030, as well as the sequence analyses presented here; five exons were identified from the coding sequence for RGS4 (approximately 8.5 kb); the critical RGS domain is encoded by exons 3 to 5; the SNPs that were analyzed are listed in the top panel; * (a small star) indicates SNPs identified by re-sequencing the RGS4 gene and * (a large star) indicates SNPs used for association analysis.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

[46] The present invention focuses on the genetic underpinnings of schizophrenia. In the first phase of the research, cDNA microarrays were used to investigate potential alterations in transcript expression in six pairs of schizophrenic subjects. RGS4 was determined to be the most significantly and consistently changed transcript. *In situ* hybridization was also used to verify the microarray findings and to examine the regional and disease-related specificity of this change. Out of the several hundred genes on locus 1q21-22, the present studies indicate that RGS4 is a strong candidate for a major susceptibility gene on this locus. Genetic association and linkage studies were conducted using two samples independently in Pittsburgh and by the NIMH Collaborative Genetics Initiative. Using the Transmission Disequilibrium Test (TDT), significant transmission distortion was observed in both samples, albeit with different haplotypes. In support of the TDT results, increased sharing of alleles, identical by descent was observed for polymorphisms in this region among affected siblings of the NIMH cases, though associations were not observed when the cases were compared to a limited number of population-based controls. These analyses are consistent with the possibility that inheritable polymorphisms in the flanking untranslated regions (UTR) of the RGS4 gene confer susceptibility to schizophrenia.

Expression Studies

[47] Two groups of human subjects, consisting of six and five pairs of schizophrenic and control subjects, were used in the present studies. Subject pairs were completely matched for sex (18 males and 4 females). The mean (\pm

SD) difference within pairs was 4.6 ± 3.5 years for age and 4.4 ± 2.7 hours for post mortem interval (PMI). The entire group of schizophrenic and control subjects did not differ in mean (\pm SD) age at time of death (46.5 ± 10.7 and 45.1 ± 11.5 years, respectively), PMI (19.4 ± 7.1 and 17.7 ± 5.0 hours, respectively), brain pH (6.85 ± 0.29 and 6.81 ± 0.15 , respectively), or tissue storage time at -80°C (45.4 ± 12.3 and 37.7 ± 13.1 months, respectively) when the studies initiated. Nine of the schizophrenic subjects were receiving antipsychotic medications at the time of death, five had a history of alcohol abuse or dependence, and one died by suicide. Also studied were 10 subjects with major depressive disorder (MDD), each of whom were matched to one normal control subject. The MDD subject pairs were also completed matched for sex (18 males and 2 females). The mean (S.D.) difference within pairs was 1.2 ± 1.4 years for age and 2.5 ± 2.1 hours for PMI. The depressive and control subjects did not differ in mean (\pm S.D.) age at time of death (52.7 ± 13.1 and 52.1 ± 13.1 years, respectively), PMI (14.9 ± 5.3 and 15.7 ± 5.5 hours, respectively), brain pH (6.81 ± 0.17 and 6.72 ± 0.30), or tissue storage time at -80°C (39.0 ± 17.4 and 39.9 ± 13.2 months, respectively). Two of the depressed subjects had a history of alcohol dependence, and six died by suicide. Two of the control subjects had also been matched to subjects with schizophrenia (685c, 604c). Consensus DSM-III-R diagnoses were made for all subjects using data from clinical records, toxicology studies, and structured interviews with surviving relatives.

RGS4 transcript analysis

[48] A Human Multiple Tissue Northern Blot (Clontech) and a ³²P-labeled cDNA probe were used to confirm the size of the RGS4 transcript reported previously (Druey, et al., 1996). However, our results reported the presence of single dark bands of ~ 3 kB in lanes from multiple brain regions (whole cerebral cortex, frontal pole, occipital pole, temporal lobe), with much fainter or absent bands observed in lanes from other brain regions (cerebellum, medulla, spinal cord, putamen). Because the UniGene entry for the RGS4 cDNA (U27768) contained only the truncated transcript (800 bp), we designed custom PCR primers based on the BAC clone sequence containing the RGS4 gene (NT_022030) to rapidly obtain the full-length RGS4 transcript sequence. For this analysis, mRNA from a control human brain was purified, DNased, and re-purified prior to first strand cDNA synthesis using Superscript II (Gibco) with an oligo dT primer. The resulting cDNA-mRNA mixture was diluted and used in a standard PCR reaction using AmpliTaq Gold (see above). All reaction products yielded single bright bands on 2% agarose/ethidium bromide-stained gels, and were subsequently purified and sequenced. Alignment of these sequences produced >99% identity matches with the BAC clone sequence containing RGS4. The 3' UTR for RGS4 obtained in this manner also aligned >99% with a cDNA entry (AL137433.1) that contains both a poly A signal and a poly A attachment site, confirming that the human RGS4 transcript is 2949 bp without the poly A tail and includes a cDNA entry not previously associated with the human transcript in the NCBI database (see below; FIG. 6).

Microarray experiments

[49] Fresh-frozen human tissue was obtained from the University of Pittsburgh's Center for the Neuroscience of Mental Disorders Brain Bank. Area 9 from the right hemisphere was identified and isolated and sectioned into tubes at -24°C as described previously by Glantz, L.A. and Lewis, D.A. in Arch Gen Psychiatry 54: 943-952, 2000, which is herein incorporated by reference. Total RNA and mRNA were isolated according to manufacturer's instructions using Promega (Madison, WI) kit #Z5110, RNAgents® Total RNA Isolation System and Qiagen (Valencia, CA) kit #70022, Oligotex mRNA Kits, respectively. The volume was adjusted using Microcon columns YM-30 #42409 to 50 ng/μl. The quality and purity of the mRNA used in the reverse transcription labeling reactions was evaluated by size distribution on a 1% non-denaturing agarose gel (>50% of mRNA smear over 1 kb; integrity of rRNA bands) and optical density (OD) measurements (260/280 > 1.80), respectively.

Sample labeling, microarrays, hybridization, and data analysis

[50] Labeling was performed at Incyte Genomics, Inc. (Fremont, CA). Two hundred nanograms of mRNA was reverse transcribed using cy3- or cy5-labeled fluorescent primers; appropriate matched control and schizophrenic sample pairs were combined, and hybridized onto the same UniGEM-V cDNA microarray. Each UniGEM-V array contained over 7,000 unique and sequence-verified cDNA elements mapped to 6,794 UniGene *Homo sapiens* annotated clusters found at the following NIH website: "<http://www.ncbi.nlm.nih.gov/UniGene/index.html>". Hybridization and washing was performed using proprietary Incyte protocols. If a gene or expressed

sequence tag (EST) was differentially expressed, the cDNA feature on the array bound more of the labeled probe from one sample than the other, producing either a greater cy3 or cy5 signal intensity. The microarrays were scanned under cy3-cy5 dual fluorescence, and the resulting images were analyzed for signal intensity. If the cy3 vs. cy5 signal intensity was within three fold, and the microarray detected spiked-in control standard less abundant than 1 copy in 50,000, the raw data were exported to a local SQL server database. On the server, the data were further analyzed using GemTools (Incyte's proprietary software) and MS-Excel 2000. Note that the operators performing the labeling, hybridization, scanning, and signal analysis were blind to the specific category to which each sample belonged.

Gene expression criteria

- [51] A gene was considered to be expressed if the DNA sample was successfully amplified by PCR, produced signal from at least 40% of the spot surface, and had a signal/background ratio over 5-fold for either the cy3 or cy5 probe. Based on Incyte's control hybridization studies

("http://www.incyte.com/reagents/gem/products.shtml/GEM-reproducibility.pdf") and control experiments, array data reliability and reproducibility cutoffs were established as follows:

- [52] 1. Genes were comparably expressed between the control and experimental samples if the cy3/cy5 ratio or cy5/cy3 ratio was <1.6.

[53] 2. Gene expression was changed between the two samples at the 95% confidence level (95% CL) if the cy3/cy5 or cy5/cy3 signal was 1.6-1.89.

[54] 3. Gene expression was changed between the two samples at the 99% confidence level (99% CL) if the cy3/cy5 or cy5/cy3 signal was > 1.9. In the control experiments, <0.5% of the observations fell into this category.

Gene group analysis

[55] Of the genes represented on the array, a G-protein group was created for data analysis, and included transcripts on the microarray for G-protein-coupled receptors (GPCR), heterotrimeric G-protein subunits, Ras proteins, regulator of G-protein signaling (RGS) molecules, and G-protein-dependent inward rectifying potassium channels (GIRKs), totaling 274 genes.

[56] At least two genes, RGS4 (Unigene cluster Hs 227571) and RGS5 (Unigene cluster Hs 24950) were mapped to the cytogenetic band 1q21-22. In order to determine whether there is altered expression of multiple genes mapped to this locus, a 1q21-22 group was created from genes represented on the microarray locus. The 1999 NCBI database human 1q21-22 map is represented by 70 genes on the microarray, although some of them are not expressed in the central nervous system.

RGS4 Sequences

[57] The RGS4 microarray immobilized probes sequence matched the entry in the NCBI database (accession number U27768, UniGene cluster Hs.227571). Of the 800 bp full-length mRNA, the double-stranded DNA microarray immobilized

probe was complementary to the 3' region of 571 nucleotides, as shown in FIG. 1A. The anti-sense, *in situ* hybridization probe was derived from the mRNA region spanning nucleotides 39-739, resulting in a 700 nucleotide long cRNA probe (see below). The RGS4 cDNA sequence, as determined from the complete mRNA coding sequence is listed as follows:

```

gtacgtcaa agccgaagcc acagctcctc ctgccgcatt tctttctctgc ttgcgaattc      60
caagctgtta aataagatgt gcaaagggct tgcaggtctg ccggcttctt gcttgaggag      120
tgcaaaagat atgaaacatc ggctaggttt cctgctgcaa aaatctgatt cctgtgaaca      180
caattcttcc cacaacaaga aggacaaagt ggttatttgc cagagagtga gccaaagagga      240
agtcaagaaa tgggctgaat cactggaaaa cctgattagt catgaatgtg ggctggcagc      300
tttcaaagct ttcttgaagt ctgaatatag tgaggagaat attgacttct ggatcagctg      360
tgaagagtac aagaaaatca aatcaccatc taaactaagt cccaaggcca aaaagatcta      420
taatgaattc atctcagtcc aggcaaccaa agaggtgaac ctggattctt gcaccagggg      480
agagacaagc cggaacatgc tagagcctac aataacctgc tttgatgagg ccagaagaa      540
gattttcaac ctgatggaga aggattccta ccgccgcttc ctcaagtctc gattctatct      600
tgatttggtc aacctgtcca gctgtggggc agaaaagcag aaaggagcca agagttcagc      660
agactgtgct tccctgggtcc ctcaagtgtc ctaattctca cctgaaggca gagggatgaa      720
atgccaagac tctatgctct ggaaaacctg aggccaaata ttgatctgta ttaagctcca      780
gtgctttatc cacattgtag cctaatatc atgtgcctg ccatgtgtga gtcacttcta      840
cgcataaact agatatagct tttggtgttt gagtgttcat cagggtggga cccattcca      900
gtccaatttt cctaagtttc tttgaggggt ccatggggagc aaatatctaa ataatggcct      960
ggtaggtctg gattttcaaa gattgttggc agtttctctc tcccaacagt tttacctcg      1020
gatggttggt tagtgcattg cacatgacat ccacatgcac atgtattctg ttggccagca      1080
cgttctccag actctagatg tttagatgag gttgagctat gatatgtgct tgtgtgtatg      1140
tctatgtgta tatattatat atacattaga cacacatata cattatttct gtatatagat      1200
gtctgtgtat acatatgtat gtgtgagtg atgtatacac acacacacac acacacacac      1260

```

acacttttgc aagagtgatg ggaaagaccc taggtgctca taactagagt atgtgtatgt	1320
acttacatgg gtgttttgat ctctgttctt tcatactaca tttgaacagg gcaaaatgaa	1380
ctaactgcca tgtaggctaa gaaagaaatg ctaacctgtg gaaagttagt tttgtaaaat	1440
tccatggatc ttgctggaga agcatccaag gaacttcatg cttgatttga ccaactgacag	1500
cctccacett gagcactatt ctaaggagca aataccttag ctcccttgag ctggttttct	1560
ctgatggcac ttttgagctc ctaagctgcc agccttcctt tcttttcttg ggtgctcagg	1620
gcatgcttat tagcagctgg gttggtatgg agttggcaga caggatgttc aacttaatga	1680
agaaatacag ctaaggcctt gccagcaaca cctgccgtaa gttactggct gaggtagggc	1740
atagaagtta aaggttactg tttttatcct ctatcctttt ttcttttctt gatcaagggtg	1800
ctcttctcat tttttctga gaaccttagc catcagatga ggctccttag tttattgtgg	1860
ttggttgttt tttctttata atggctctgg gctatatgcc tatatttata aaccagcagc	1920
aggggaaaga ttatatttta taagagggaa caaattttca caatttgaaa agcccacata	1980
agttttctct ttttaaggtag aatcttggtt atttcattcc aaacatcggg gctaacagag	2040
actggaggca tttcttttta ggctctgaga ctaaatgaga ggaaaagaaa agaaaaaaaa	2100
aatgattgtc taaccaattg tgagaattac tgtttgaaac ttttcaaggc acattgaaat	2160
acttgaaaac ttctcattta tgttatttat gatgttattt tgtacgtgtt attattatta	2220
tattgtttta taaatggagg tacaggatat cacctgaatt attaataaat gccaggaag	2280
taattttctt ctcatcttc taaaactact gcctttcaaa gtgcacacac acgcgtccac	2340
atacactgca ttcgttgctc cagtataaat tacatgcatg agcaccttct tggtttttaa	2400
gccaatataa tgggctgcaa aatgaagaca ccagagtgtg tgcatacaaa tctcactgta	2460
ttaaagatgc aggttttcta attgtaccct tcttgtctct ctggcaatct tgcccttaat	2520
atccctggag ttctcatca gtgtcatttt ctgttataca cagttccaca attttgtctc	2580
tagttgactt caaatgtgta actttattgg tcttgccta ttataattgt catgacttct	2640
agattgtatc tgaactcaca gactgctgtc ttactaatag gtctggaagg tcacgctgaa	2700
tgagaagtaa attattttat gtaatacatt tttgagtgtg tttttcagtt gtatttcctt	2760
gttatttcat cactatttcc aatggtgagc ttgctgctc atgctcctg gacagaatac	2820
tccttccttt tgcatgcctg tttctatcat gtgcttgata ggctcaaag ctaatgcttc	2880

cagtgaaca cacgcatctt aataataagg gtaaataaac gctccatg aaac

2934

[58] For purposes of the present invention, the RGS4 cDNA will be referred to as SEQ ID NO:1.

[59] The 205 amino acid long sequence of RGS4, as determined and reported by Druey *et al.* in Nature, 379: 742-746 (1996) which is hereby incorporated by reference in its entirety, is listed as GenBank Accession number P49798 as follows:

[60] MCKGLAGLPA SCLRSAKDMK HRLGFLLQKS DSCEHNSSHN KKDKVVICQR
VSQEEVKKWA ESLENLISHE CGLAAFKAFK KSEYSEENID FWISCEEYKK
IKSPSKLSPK AKKIYNEFIS VQATKEVNLD SCTREETSRN MLEPTITCFD
EAQKKIFNLM EKDSYRRFLK SRFYLDLVNP SSCGAEKQKG AKSSADCASL
VPQCA

[61] The above amino acid sequence of RGS4 is referred to as SEQ ID NO: 2 for purposes of the present invention.

[62] Untranslated regions upstream and downstream from the RGS4 coding region are identified in the context of the present invention as being relevant components of the RGS4 gene. The RGS4 coding sequence along with these sequences are found on NT_022030 as described in greater detail below. This sequence is

agttcaagac cagcctgagc aacatggtga aaccccatct ctactaaaaa tacaaaatta	60
gacagggcatg gtgatacacg cctgtaatcc cagctacttc ggaggccgag gcaggagaat	120
cacttgaacc tgctgggggt ggaggttgcg gggagcaaga tcatgccatt gcactccagc	180
ccaggcaaca agagcgaaat gtcattctcag aaaaaaaaaa aggcatttta tatatatata	240
tatatatata tacacacaca cacacatata tatatacaca tatatatata catatatata	300
tatatacaca tatatacaca tatatatata catacatatg tacacatata tatacacata	360
tgtatacaca tatatacaca tatatacaca catatatata catatatata cacatatata	420

cacatatata cacatatata cacatatata catatatata catatatata tatatacaca	480
tatatataat atacacacat atatatacac atatatacac acatatatac acatatatac	540
acatatatat acacatatat acacatatat acatatatac acatatatat acatatatac	600
acatatatac atatatacac atatatacat atatacacac atatatacac atacatatac	660
acacacatag atatacatat atatacacat atatatacgt atatatatgt atatatatat	720
gctccagagt tcataagagg tagcagttga ttaccactgg ggatagagga aaagagagtt	780
tgacagcagt gtattgtgag aaggacattt cagggttgatg gcaaatagta ggggaaatac	840
ataaatgtgt aataaaacct atctgtaagg tagttaagaa ggtaaacacta tatatatata	900
tagtgaaagc agtgtaaacc taaaggatgg gccaaaggatt taaatgttat agaagaatgg	960
ctaagatgcc aaagctcagt gtatgtggca gaggcattgt gtaggggtgtg tccaggttca	1020
tatattgcat taagtgtgag aacaccctgg agtatgaacc aagaaaatgc aaaagccaga	1080
agtgatggag gaaatgagac acaataatga agatattgag aggaggggtgt gggcctagag	1140
tgaagctttt cgtgccagta cttcttttga agggccagtt ctcttctctc tcgggggctc	1200
cttcattctc catagagtcc acagctttta agggccaaca cttgagggtca gcctggctct	1260
ctcatttgag ctggatagaa cattttagag caccatctat tcttcaagag gaagtttaaa	1320
aataaaagaa ccttgaagag gaaaaaatgt agacattcaa tctaaccctt tcattttact	1380
agccaaagct aaatagaatg caggttacct gtttttcagc caggcaccat catttcctaa	1440
ttgttataaa atttattatt attgttggtt ttattattat ttgccataag aagtttccca	1500
tatcctttta gtataacaaa aacacaattc acaagcatta taaaacccat ggtgtctaac	1560
tattaaaaaa attaagtga acacacttgt cccagctact ggggaggctg aggaggagg	1620
atcacgtgat cccagggggt caaggttatg gagagctatg attgtgccac tgcactccag	1680
cctgggtgac agggaaagac cctgtctcta aaattttttt taaaaaaact aaactggttt	1740
tattacagag attctggaga cagctacaca taaaaggggtg gtatgcctca tattagctac	1800
ccaggagggt ggaatgccaa cttagggtgt gtcaccacta ttaaaaatgc cccaaagcaa	1860
tcaaaactga gaacttctg ggagcttagc attgtgcaaa agcagcacia aacacttaaa	1920
caattcacag ttgtgttga atgggaaggc ctggaaatat aaaccaaaga gtatattgtc	1980
taaattgata gagattacaa ttgcctgaaa gaaaaagttg actttttaact agaatgttca	2040

gagtaggttt acagaagaag ctcttaaact gggctccagt ggatttgta atgctttgga	2100
agctgggtggg gtgggagggg tggagggggc ataaaaagtc atgttggtat gctctgctca	2160
agtctccatt ctgtttcctt ttctcttttt caatgtcatg tcccattatt tcattatggg	2220
cttcccttta tccaggatca atatgccacc tcttggttgt cttttaccta cttctccacc	2280
tcactatgga atcgtccttg ggtagctcct gtgcttgga accctgcacgg gcacttttct	2340
gatgtcttga ttccagcttt actcctaaaa cttaaagtct gaggggcca caccatggca	2400
gtggtaggga tgggaatggg ggtcttgtaa cacactacat aaactacacg aaataaacta	2460
catgaaactc aacatgtttg caagactcag ttcacatcca tgaggagctc atgcttctcc	2520
ctctgctcc cctagcacac atgattatct ctatttgga atgtttgga tttttggtga	2580
agtgaatggt tcaataactt tctccaccat cagaacaaaa gctctttaag gttagggatg	2640
ggatcataca cacttccctt gtccaagtcc ccatcacccc ttatctagac aattgctaca	2700
gtttcctaca cactcttcta acctcttgca gtctattttc ataaaacagc tagagaactt	2760
tgagatgtaa gtcaaaaaat agaacatgtc gctctttccc attgtttttg aaataaagtt	2820
caacccctt accaggggtca acaaggccct gcaatgattt ggctctgtta aaaattcttt	2880
agccttaact catgctgttc ttcttacac tcaactgcatt ctagccattg aggtttctat	2940
gcatcaaact ttttttggtc ccagcactgt gcacatcctt ctgggtagaa tgccccttga	3000
tttgataat tagcacctcc ttcacatctt aggtcttagt ataactacta ccttcttaga	3060
gaagctctgc ttcttcatcc tataaaaaag taaaattcct taccctgtta ttttttaagt	3120
catccgtgtt tcattctgtt aaagtctta tcacaattta tcattatttt atttacagtc	3180
atgtgccaca taacaatggt tcagtcaggg atagaacaca aatgtatctg gcccataat	3240
attataagct gagaaatttc tattaactag tgatatcgca gccatcataa gtgtaatgca	3300
ggacattacc ttttctatgt ttgatatgt tagatacaca aatatatttc attgtgttat	3360
aatttcctac agtattcagt acagtaacat gctgtacagg tttgtaacct aggagtaata	3420
ggctatacca tacagcttag gtgtgtagta ggctataacc atctagggtt gtgtaagtac	3480
attctatgat attcccacaa tgatgaaatc acctaacac acatttctca gaatgtttca	3540
ctgttgtaga gtgacccatg actatatctt cctatatact tgatatcttt gtgcatctgc	3600
ccatgagaat gtagtgtaag atcaaaggat gcaagaatgg gttctatcca gtatagtacc	3660

cactacactg gtggatgtca atatgtatTT gttagattaa tatctcaaga atgagcacct 3720
 ttctcagaca cataaaagat gctcaatata aaagtttgTT gaactgaacg ttattggcaa 3780
 atgtaacatg atcggattta aagaggagcg aaacagaggt ctggctcaaa caccatactt 3840
 ctagagtgca taagaggtag cagttgatta ccactggcga caggagaaaa aagagcttga 3900
 ccgcagggta ctgtgaagac atttcaggtt gatggcacag aacaggggaa atacataaat 3960
 gtgtgggaat attcagtggT ctgggatgac tacatagtag aatataatga agaaaagagt 4020
 ggaagggaaa gatgaaaagt tggaatgggg atgaattatg aaagtaccag aatgttatgc 4080
 taaggaatct agattttaaa atgtgagggc aaattgaagt **C**ctgggcacg ttacaaaact 4140
 agaggtcata aagtttacc taattacca agatttccta gaggatctat aattggaatc 4200
 cagatctgcc tctctgtaaa gttcaagcac ttccatgac accatactgt ttctttccac 4260
 ctgcacaatg caaatgaact cttatgaaac tgctgtttct atcctgggct aaatgttgca 4320
 gaaaaaagat ttaatctttg ggataaggct attttgggtt ttctcct**a**ct tcttgggaaa 4380
 caaggttttc ttccctggc taattaagtg tggattgtt cttccagga aatcagtgat 4440
 gcatcacctg ctgctatcaa atgtcagggT tggagtTcct gatttattgc atgtgccac 4500
 aaagcttggT gcaaagaatt ggacacattt cccaaaagta agacatactg ggaagtcct 4560
 gtttaccttc ctggtataca gcatcctcca gccccatctt tttgcttttt agtcctaaaa 4620
atcaataact gaactctcat tgatgtctag gccattgtag taaacaataa agaaggaggg 4680
 aggttctga caactgagag gaaattgtca tctgaagtgg tgcaagcaca gcctggggct 4740
 gagccttggc ctacatcctg cccaagtggg ggatcagtg**C** ccatttaac atctggtaga 4800
 actaaagaac gcaac**g**cctg ccacaatgac ttatttcctt gcatttgata ccgtcaatcc 4860
 ttgagaaatg tttctttttg ttctccctga gcaaagggtg gaaaaatttg aaatttacct 4920
 agagaccaca catagttcac atcctgctgt gtggctgaat gtctgcccc**C** cagtaggaaa 4980
 cagttcttct aaagcctatt gtcaacaata ccttcagat gttagcattt tacaatttaa 5040
 ggaacttaaa atag**C**ttca aactttttgc cagtttctct gatatccaat ctattctttt 5100
 actctgctc ccaagctttc ttctagaat gctaacctga tcggcttaag tacttgaact 5160

acctcttctc ctccattaac tacagagtaa attctggtct tcagagtaac aagaaacacc 5220
ctttagttct cagcatattc gtgcaccttc atttatctct ccttctctct caaagctgca 5280
gtaggggtga aaac**g**tgtga tacatcttct cttccatcat aagggtcgca accaaaactc 5340
ctatagtaaa agacaggtta ataagagcaa aacctaacaa atttatttaa tcaaagtttt 5400
acatgacatg ggagtcttca gaaatgaaga ccaaagacc caggggaaac tgtctgtttt 5460
ttttgctgag gttcgatgaa gaatggatag catgtagcca tgtagattag acaaaggat 5520
atgatctagt ggtaaaggac tcagggggaa acacagcaag gcctgtctat tcagattctt 5580
cttgatctct ctctctctat gtatagcatt ctttctctct gagtatgggg caggactctt 5640
cttcaatgag ggtcttcaag ggagaaggga gaaagtggcc tttttagatt ttat**gg**cttg 5700
cttcggggaa gaggagtctt agtttctatg acccatcttg gggaagagga attctggttt 5760
ctgtgacttg ctttcatgaa gaaagaggag taagaggcag gagggcagga gatggtcaga 5820
aagagacttg gctgcttctg agggcttccg ctctccttta gttccaagta cttcttagca 5880
taccaaagca ctatacttg gcatatggtt ttctgagctc taacactgca atcatgctaa 5940
actctctat gaccttcaaa cattccactt gcttttattc tttatggttg tgatggcata 6000
gaggtcaata gcaaagaccc tggagtccca ctgtctgagc tggcataaca ttactaccac 6060
ttaatcaatg tgtaagctca ggtaagtact taagtctct atgcttcato tgtaaaatga 6120
gaatcattga agaacattct ctcaggatgg atcatgagga ataagtgaat taactggcat 6180
atagtgtta aaccagtgcc ttgctcagtt agtgacagat aaaatcatct gttattactg 6240
tgcccactat tgtgatgctc ttctcttctt tgtacaacga ctacatctct atttatcatt 6300
ttaggggtctc cttgtgaaaa accactccag attcaaaaga ttgagtttaa tctctatctt 6360
ctgtgcttct ctggagtttt gtaaagtaaa tcttcaactg acatcatgga taggttcttg 6420
gaaactacaa cttcaagtga aaggacataa ctaaaccaat ttttttctca tcaacgttat 6480
aatgaaatgg cattgatgaa atgatggcat tcaaggacct gctgtacctt gtttcaacta 6540
aagtcactgt ttccaataat ctattgatga cattgaggac ttactatata ataataaata 6600
tatatataat cgacgaaaca ggaatcaaac tgctaactct gctaactggc ctcctgctt 6660
ccacactctg cccactcatc tcagtcttct tttcacaaga gtcagaatga tcagatgaga 6720
ccctctctct gcttctgttt cttccatgga tttccactgc actctgataa agtccagcct 6780

cttgaccaca gectacaaat ccttgcacga tctatcgttt acttttccat ctctttttat	6840
gctactttca tcttgttctc aattctctag ctatgctggc cccttcttgt tctttcccat	6900
ttttttttaa tttttaaaat ttgtatatat ttatgggtta taagtgaat ctttttagat	6960
gcatagggtg tatagtata aatcagggc ttttagggta ttcacacct gaatgatgta	7020
cattgtaccc cttaagtaat ttctcaccat ccgctgactt cttgccccct gggatttcat	7080
cacctgaatg atgtgcattg tacccttaa gtaatttctc accatccgct gacttcttgc	7140
cccctgggta ttcacacct gaatgatgtg cattgtaccc cttaagtaat ttctcaccat	7200
ccgctgactt cttgccccct catccttctg aggtccatt gtccatcatt ccacactcta	7260
catctatgtg tacacattat ttagctccta cttataagtg ataacatgca atatttgtct	7320
ttctgtgtct gtcttgttt acttatgata atggccccca gttctatcta ggctgctgca	7380
aaaggcatga ttctattctt ttttatggct atgttcttct ccaatttaga taaagaacac	7440
tcgcacttgc tcttacttct atttgaata ctaattccta ggcttcttgc attgctttct	7500
ccttctcacc catcaaactt catttttagat accacctctt caaagagggc tttcctgacc	7560
accttggtg aattagccct tcaccatctg attactctct agcacatcac ctgcccattt	7620
tattcatggg acaggtcaaa atctggaatc acctgatttg tttattttct gactccttct	7680
actgagatga aaactctact agagcggaga ttttatctgc ttgtatcagg tactgcttca	7740
aacagcacct gatacaga ^{gt} aggtgggtcaa aagatatttc ttaaacaat gaacaaataa	7800
aaagtagatc ttttgagagt aaagctcttc cacactacca gagtcattca ggaatgacaa	7860
atcatagaat aacagaattt gatgctttgt gcatatcaga gaaagaagg ggaagggtgt	7920
caaggatatca tgatgtacca gtccctgcct cctcaaacac aatctgcaag tcccacagt	7980
aaaaagtaag ttaactcatg tgaagcggtt tacaacact tttttaaaag tcttaaaact	8040
cctaagaaag caagatttaa tagtcaaaga agtgagtaaa catgaaatgc ctgaacagag	8100
taatgagcta agcacaaagt tagagacatg ttagttaata tgtcttgaaa gcagcagctc	8160
ctgctttcaa ggagcaagaa caaattgggc aagtgaacac tccttgaata aaatgtgtaa	8220
aattaatttt gggttatgtt ctatactgtg tataatagaa tgataaaaat tatttgacta	8280
gcactttgta gtttagaaat atctctattt acacagttta ccttatttga taagactggt	8340

gagtgatggg atagcatggt ggacaatcca cataactgag tatcgagaca cctgtatctg	8400
gaccagctc tgtagtaag aagctgtaac ctcagcaagt cactttctct ttctgggtct	8460
ctatttcctt tttggtgaaa tgagagtgtt aggctagatt gcctttgaag tccatttttg	8520
tctttaaagt cccatctatt gcagtgattt atatttaact catgacaaat caggcttctc	8580
ttattctaag tgcaa g acat aaaactttta ttgtggaatt tcaggcatca gtaaactctt	8640
ttgggtactc acttatgttc ctgaaatcaa tctatttgag tgatcactct tttaggtgcc	8700
caggtaaaca aagaaggcca tggctcttct ttgagtgacc ttctttccct ttttaattagt	8760
ctgacctctt taatgtcagt tctgactgat tcatttcctt ggtccatctt ccttggtctg	8820
agggccttcc tagtttcata ttgcacttca gttccttcca caccaccatc aaggatggct	8880
gtcaacattc atttgttcta tggtataatt caaggaaaag ttgccagta gctaatacaa	8940
taaatgcctt cttatgggcg gctagagact ttttctata atttaaagtc atcttctgta	9000
gattatggtc cctccaccac ttacatttg tctgctgtct ccttgctctg ctagtcatgg	9060
aacgtgttgg tagtgggggc agtgtgggat gttcaagggc acgtattggg tagggccaca	9120
tatgggcatt gctttgtgcc attctttcta tatttttggt attttgcac tcaactggaac	9180
ccaactattt ttcactctt ccacctaaac tatttgatgc ctctgtttct tataataaaa	9240
gtatagctca ctgtagccta tgatcaggaa cctatctgct ttctaaatga aagctgtttt	9300
ggtcagatct agcaattaat tcccttcttc cacttatagc tttcctctgt aactctggtg	9360
taggtatttg gtttatggct ataagatgtg aaacacctga atgattctgt ccatgcaggc	9420
atttcagttc atgatattgt atgtaaaaga tactgattgt ctaggtgttc agaaacacct	9480
atagggctta atattcttac aatcagtttg aaggctgggtg atacgcaaag caaactacat	9540
atttttctgc ctgctctctc tctttctctc tacatctctc tttctttatc ttttgaaata	9600
tcagtttgga gacttagaat tacataagac ataaacccat ttgatataag aattgctgtg	9660
tatatttgct catctactcc ctcttttggt cctcgagctg ccggtttaga ctttttacag	9720
gacgcaggca tgtgaaggag aaactgtcag tgctaggtg aattctgttg ttaccaagat	9780
ttctagaaaa gtattcctca gtcaggttga ttacagatat agcaaata tttttcctag	9840
ggtagtttct gtatgctgcc gggcttataa ctgtctgtca tccagctatt t C ctctccacc	9900
ttcttgtttg cataacaacc aaggcaactt ccgcaaatca ctgcgtggag acgatgatcc	9960

tg**C**cagctcc cttttggaaa tegtgaggat cagatcttgg accatgtata atatgatget 10020
 tctaataccaa aagaggaaaag gcattgggag tcagctccta agtaagctcc agaattcctg 10080
 ctggtactttt tccttccagg aagcaacttc cttgatattt tttttttaca gg**g**catatgaa 10140
 taaaaactat attttgcagc attgtacact ttttttcctt ttctagaaat tctaaacctc 10200
 tgacattggg ggagacattg agtacatttt ttcccatatc cctacttttc agaaggattt 10260
 tctctgctcg ttcacttaac attgctgatg cgtcagcttt ttcttctca tctctttcag 10320
 gggctggaga ggcagagga gacagaggag ctggtactgc agagcggtcg tctgattggc 10380
 tggacggtcg tagctgggct ataaaagaga cccctacagg cttagcagga agacgctcag 10440
 aggattctga caatatcttt accggagaag aggcaaagta cgtcaaagc cgaagccaca 10500
 gctcctcctg ccgcatttct ttctgcttg cgaattccaa gctgttaa ataatgtgca 10560
 aagggttgc aggtctgccg gcttcttgct tgaggaggta agattgcttt cagccattaa 10620
 ccatattaaa cttttggcta gactttctca gttatttaca tgttgactt actaacctag 10680
 ttctgtgcaa ttagaaacag tgtggtcagg agagcacgac tttctaactt tctccaaga 10740
 ctagctagat attgtgactt aagacatgtg ctcccaa at ttcagccctt atgtgttggt 10800
 ttgtgtgacc tcagttttga gaactgttct attctttaag ccaggcttaa gaaagctagt 10860
 ttttaattaag aagcgagatg aggtttgagg ctatgtacag tgatctgtaa tatctccatc 10920
 tgtgattact actgctattt gagcatccct ggagtacata gaagcctggc tctgggcttt 10980
 ctgattgtat gctacaactt gtttcaggaa aggtacccca gaatgagggt tggtccatc 11040
 atcagaaagg cacta**t**gctt tccgtgtggt ggtgcagtaa ctttactct **C**tatgttctt 11100
 ataag**C**aaat gttacaatga gatatgagtt ttaaagccag atcttctta tctctctgcc 11160
 ccactcttag ttcttgaagt gtctcatatg agtttggttg agaaatattg atcattacaa 11220
 atcagttaat agttttgtag aagatctcat cttaaagaca ttgttttggt aatatactcc 11280
 cttgattttt ttaaaagacc ttacagacat acagctattc atttgttttt gggttggtca 11340
 aaaaaggat aaagaaatgc attcagagaa agatcatata ttagccagtt gaaaattaaa 11400
 cacaaaatga gtgcatatta cactacttaa tcttgcagtc aaaggtaaaa agtcaacct 11460
 aaggatatact acctgctttc ttatcgact gcaaatagaa attaccacaa attttatttt 11520

ggaaataatc tcagaaaaca taatttttta tgtactatta aaacatttac tttccaaata 11580
 ttctgtcatt caggagtatg gaagtatcga tggcttcttt aaaatgaagc aggaggggtct 11640
 ggcagagagt atctatgaaa taagttcctc tgaccttcac gcttaatttt ctgaatggag 11700
 tggagcaa at tacttcaagc ttcacttaac ttgcatatga aatgaaccgt acaaaaatac 11760
 aagagtgtca gga**g**aaagtt atgctctggg aaatattttg caaaacagat aaaagataat 11820
 actagagctc tgtcctcaaa gagttaagca gctaatactaa ggaggtaa ac tctatgtcag 11880
 caggatgaac tgctcttccc tttcctcctc aataaattgc aaatcatcta gtccaacatc 11940
 tttaccacca gtgcctgagg ctccagagga gccattgcct tctcaaggtc acatagggtg 12000
 tgggtgagtt aggaccaa at ctagaattcc tgactccagt aacttctgaa gtcattttgt 12060
 tttttatttt tatgggtttta ttataagaat acttgctaag cacacttacc ccttgcattg 12120
 attaataact ctaggatctc ag**gtg**gatcc agcacataga aatatgaatt cgtttctatt 12180
 tggacttcat gatatattta cattatcacc ttggaatcac cctaacattc aggattgtat 12240
 cttgttataa tcaaaaagga tggtgcatcc cctgaacagt catcagtcag ggaagcagag 12300
 gagggaaagt aatcttgca ggaagagaaa atactattta agggacagtc agagaacata 12360
 atggaattca aactttctgg gaaaacctac atacataaat gtattagtgg ccatcctaaa 12420
 tgtctttata tctttgaggc tttattttcc ctactccaaa tagacacatt tagttattca 12480
 tttcttttaa aatggtat ttttttttaa actatttctt gactttttta ataaaaagag 12540
 atgcaagcaa gaggatattt aataaaaagt aagagagttg agcttaaggc ttattaaaag 12600
 accccctttt tctagttagt caggagctct aatgtgccct ggctacctat taaatgggtg 12660
 caataaactg gaagctcagt gatgactcta gcctgcttct cctaatagct gttaagcctc 12720
 aaatgccctt tagagtgtgt atgtccttta aagtagctat taagaaggaa agcagcagca 12780
 gcagatattg tctagaaaaga agccccaaga agctgagggt tcagcttggg catttgtttt 12840
 cgccatcca tgctccattt ccctctgctg gaactgtgca cctcagtgta ttctccctct 12900
 atacctcaca gcaggaactg cttgcccccc ccccccccc ccaacatata tggctggaac 12960
 tgaatagact tttactttcc cgagggtgct ctacagttcc ctctgccagc aggggaacag 13020
 atggaaatag caatcacctg ccagaagggt gcgtgcagca aggatgtgca tcttttgccg 13080

ctactgcttt ctgattccta aaaattactc agagatcact catgtgttca gtgattcagg 13140
ttctgttgaa gataccaaag atattcggtt ggtcaaaatg acgggcatat aaaggcttct 13200
caggtttctg aggtaaactg aagggtcaga attccagttg tggatgaagg aaatgggtgtt 13260
atgactgcct caaggttttg tagcaagtca tagggaacca agaggaatct tgttttcctc 13320
agaggtcattg ccaactccaa ctcccgttcc ctaaaactgtc tctgagccat agactagtaa 13380
tggactcttc aagctctacc attaggtatc ttttaaagaa agctgggttat tactatttat 13440
tcattttttt ctcttctgtg cagtgcacaaa gatatgaaac atcggctagg tttcctgctg 13500
caaaaatctg attcctgtga acacaattct tcccacaaca agaaggacaa agtgggttatt 13560
tgccagaggt aagagaaaag gccttggtga agatgtactt agtattaact atctgatgat 13620
ggggatgttc tgtgagaagg aacttggtgct cctagttaag ccagatttgg atcaagatag 13680
cctccatttt catggagatc ataactacat ttgaaatttc tatacattta gtgaaaaact 13740
gccctcatca ataacatatt ttgtcataac gatggaaaat aaaatctttg ccttcattca 13800
ggatcttaga tttcttgccc caattttttt accatggcat tccaattatt ctgtttctct 13860
ctattttttc tagagtgagc caagaggaag tcaagaaatg ggctgaatca ctggaaaacc 13920
tgattagtca tgaatgtaag tctgacagca acctgggatg aggtactctg gataagacaa 13980
gttatattat gctgggtctaa tagaaactgc agcaaggcct ggcttctttc tgatgttcag 14040
actcaggaga ctcttttaggt cttaaattca gtctgtttta aattttaata tgccctagag 14100
ctttgtgata tacaatgaaa agtttatgca ggaaccatgt ggaaaaccat ctctctcatc 14160
acaaggaaaa acggaagaga gaaaaaaaaat gataaatatc aataccttct tgcaaaatca 14220
atctcagttt ctctttccca aattgacctt ggtaattgat agctgcatag gcatttcaga 14280
agcaaaatac ttccttgaaa gaggttcca acttgagtaa gaatcattag gtagaactgg 14340
gaaccactgg atatcaaca cagattaggg ttacctgact ccagggtgact tgaaaaaagc 14400
aggggaaaaa gggattgctt gaatccatgc tttatcccc aagtacctca gctttatgtg 14460
aaatagcata tccaagaggc caaccagtgt gatgacaact gtggtccttt ctctgtatc 14520
ataggtgggc tggcagcttt caaagctttc ttgaagtctg aatatagtga ggagaatatt 14580
gacttctgga tcagctgtga agagtacaag aaaatcaa atcaccatctaa actaagtccc 14640
aaggccaaaa agatctataa tgaattcatc tcagtccagg caaccaaaaga ggtaggtttt 14700

ttatggatac	ataaaaattg	tacgtattta	tggagtatgt	gtgatatttt	gatacatgca	14760
tacaatgtga	taacaatcaa	atcagggcaa	ttgctatata	catatctcaa	acattttatta	14820
tttctacgtg	ttgagaacat	tccaaatctc	ctcttctagc	tatcttaaaa	tatacaataa	14880
actattgata	actatatcac	cctaattgtc	tatcaaacac	tagaacctat	tccctctacc	14940
caactttcta	tctattcctt	ctaccatta	gccaacctga	ccaaaaaggt	aagcttttat	15000
ggcagagaac	tctctggatc	ttagtgaagg	ttcctagaat	agtggagctg	actatcataa	15060
tcttgacaac	cccaaataaa	tcagtttttt	aaaaaatctc	ttttatccat	gtggcttacc	15120
ataacctccc	tgcatgaatt	tttctgatga	atctccccaa	tttgtagac	agaacagaag	15180
atcttgccct	gctctctcta	aagcagaaa	gttcattctg	aacctttcat	actctctcac	15240
atgtgccaa	gaggacccca	atgtcacttt	tgttttttgc	ttctgaaata	cagaggggtgc	15300
actgccactt	acaagtcact	acaaagcata	caggcttgca	tcctcaacag	ggatataggt	15360
ctaatgaagc	cttggccttt	gcccctcagg	tgaacctgga	ttcttgacc	agggaagaga	15420
caagccggaa	catgctagag	cctacaataa	cctgctttga	tgaggcccag	aagaagattt	15480
tcaacctgat	ggagaaggat	tcctaccgcc	gcttctcaa	gtctcgattc	tatcttgatt	15540
tggccaaccc	gtccagctgt	ggggcagaaa	agcagaaagg	agccaagagt	tcagcagact	15600
gtgcttcctt	ggccctcag	tgtgccta	tctcacctga	aggcagaggg	atgaaatgcc	15660
aagactctat	gctctggaaa	acctgaggcc	aaatattgat	ctgtattaag	ctccagtgtt	15720
ttatccacat	tgtagcctaa	tattcatgct	gcctgccatg	tgtgagtcac	ttctacgc	15780
aaactagata	tagcttttgg	tgtttgagt	ttcatcaggg	tgggacccca	ttccagtcca	15840
attttcctaa	gtttctttga	gggttccatg	ggagcaaata	tctaaataat	ggcctggtag	15900
gtctggattt	tcaaagattg	ttggcagttt	cctcctccca	acagttttac	ctcgggatgg	15960
ttggttagt	catgtcacat	gacatccaca	tgcacatgta	ttctgttggc	cagcacgttc	16020
tccagactct	agatgtttag	atgaggttga	gctatgatat	gtgcttgtgt	gtatgtctat	16080
gtgtatatat	tatatataca	ttagacacac	atatacatta	tttctgtata	tagatgtctg	16140
tgtatacata	tgtatgtgtg	agtgtatgta	tacacacaca	cacacacaca	cacacacact	16200
tttgcaagag	tgatgggaaa	gaccctaggt	gctcataact	agagtatgtg	tatgtactta	16260
catgggtgtt	ttgatctctg	ttctttcata	ctacatttga	acagggcaaa	atgaactaac	16320

tgccatgtag gctaagaaag aaatgctaac ctgtggaaag ttggttttgt aaaattccat 16380
 ggatcttgct ggagaagcat ccaaggaact tcatgcttga tttgaccact gacagcctcc 16440
 accttgagca ctattctaag gagcaaatac cttagctccc ttgagctggt tttctctgat 16500
 ggcacttttg agctcctaag ctgccagcct tcccttcttt tcctgggtgc tcagggcatg 16560
 cttattagca gctgggttgg tatggagttg gcagacagga tgttcaactt aatgaagaaa 16620
 tacagctaag gccttgccag caacacctgc cgtaagttac tggctgagtg agggcataga 16680
 agttaaaggt tactgttttt atcctctatc cttttttcct ttctgatca aggtgctctt 16740
 ctcatTTTTT cctgagaacc ttagccatca gatgaggctc cttagtttat tgtggttggg 16800
 tgttttttct ttataatggc tctgggctat atgcctatat ttataaacca gcagcagggg 16860
 aaagattata ttttataaga ggaacaaat tttcacaatt tgaaaagccc acataagttt 16920
 tctcttttaa ggtagaatct tgttaatttc attccaaaca tcggggctaa cagagactgg 16980
 aggcatttct ttttaggctc tgagactaaa tgagaggaaa agaaaagaaa aaaaaaatga 17040
 ttgtctaacc aattgtgaga attactgttt gaaacttttc aaggcacatt gaaatacttg 17100
 aaaacttctc atttatgtta tttatgatgt tattttgtac gtgttattat tattatattg 17160
 ttttataaat ggaggtacag gatatcacct gaattattaa tgaatgccc ggaagtaatt 17220
 ttcttctcat tcttctaaaa ctactgcctt tcaaagtgc cacacacgcg tccacatata 17280
 ctgcattcgt tgctccagta taaattacat gcatgagcac ctttctggct ttttaagccaa 17340
 tataatgggc tgcaaaatga agacaccaga gtgtatgcat acaaatctca ctgtattaaa 17400
 gatgcagggt ttctaattgt acccttcttg tctctctggc aatcttgccc ttaatatccc 17460
 tggagtccct catcagtgtc attttctgtt atacacagtt ccacaatttt gtctctagtt 17520
 gacttcaaat gtgtaacttt attggtcttg cctattata attgtcatga ctttcagatt 17580
 gtatctgaac tcacagactg ctgtcttact aataggctcg gaaggcacg ctgaatgaga 17640
 agtaaattat tttatgtaat acatttttga gtgtgttttt cagttgtatt tcctgttat 17700
 ttcatacta tttccaatgg tgagcttgcc tgctcatgct cctggacag aatactcctt 17760
 ccttttgcat gcctgtttct atcatgtgct tgataggcct caaagctaat gcttccagt 17820
 aaacacacgc atcttaataa taagggtaaa taaacgctcc atatgaaact atttgcttgg 17880

aaacacatta atgatccaga gacatgctat gagaaacatc aggggtgtagg gtgacttttag 17940
 aaaaatactc atactgagtc tttaatccct cctgtgccag tgaactctgg gaaagaaagt 18000
 acaaactgaa tattgtttat tcttttagttc atgccactgc tctgcttggc tctactcata 18060
 gaaccaaggc aatcttagct tcagagactg caaaacagat taagtgattt gcttgcagat 18120
 tctcaatcaa ttttcaaggg atagagttca ccttccagag ccattctttt atttccagtt 18180
 acccgctgt ttgagagatg atagagcagt gggaaattga gagagttgaa aggagctata 18240
 gattcttacc caaacttcaa aaatccttcc ctcccttttg ttaattctct ttcctggaaa 18300
 agaggtcata aaatgttcac atcctcagta ataggccctg tgctgtgtct attatgtcat 18360
 gagactccca tttcctgacc cttctttccc attgtaagag tagtagttac aagggtgttaa 18420
 ggatagatga tcttcaacac ttttgagaaa tagatccatt tacggatctg gtaaaaaacta 18480
 tggaccgaac catcttttaa gaaaaaaatt cagagaggaa tctaaatttt gtgtgctttg 18540
 aggggaaact ctcagaatct cccctcaaaa ctatcattct tctcttatac tatagatgtg 18600
 tcagactctc actgggactg tatagttgct gctccctgta tttgataata tctatcaaga 18660
 actgcagggt aattcaaagt cacgctatta gcagcaagtg tgagcagtggt tggtttcccc 18720
 agtctctaca tccctcatcc tttctttctt ctttatgggt gtctattaaa gaaataaaaa 18780
 aaaatattgg ctgaccgttt ttctgaagat aatgtatatc aaggaccacc ttttgaaaaa 18840
 cactcattat tcgagaacaa agacacaaca tacgagaatc tctgggatac attcaaagca 18900
 gtgtgtagag ggaaatttat agcactaaat gccacaaga gaaagcagga aagatctaaa 18960
 attgataccc taacatcaca attaaaagaa ctagaaaagc aagagcaaac acattcaaaa 19020
 gctagcagaa gacaagaaat aactaagatc agagcagaac tgaaggaaat agagacacaa 19080
 aaaacccttc aaaaaattaa tgaatccagg agctggtttt ttgaaaagat taacaaaatt 19140
 gatagactgc tagcaagact aataaagaag aaaagagaga agaatcaaat agacacaata 19200
 aaaaatgata aaggggatat caccaccgat cccacagaaa tacaaactac catcagagaa 19260
 tactataaac acctctacgc aaataaacta gaaaatctag aagaaatgga taaattcctc 19320
 gatacataca ccttccaag accaaaccag gaagaagttg aatctctgaa tagaccaata 19380
 acaggctctg aaattgaggc aataatcaat agcttaccac ccaaaaaaag tccaggacca 19440
 gatggattca cagctgaatt ctaccagacg tacaaagagg agctgggtacc attccttctg 19500

aaactattcc aatcaataga aaaagagggga atcctcccta actcatttta tgaggccagc 19560
 atcatcctga taccaaagcc tggcagagac acaacaaaaa aagagaattt tagaccaata 19620
 tccttgatga acattgatgc aaaaatcctc aataaaatac tggcaaaccg aatccagcag 19680
 cacatcaaaa agcttatcca ccatgatcaa gtgggtttca tccctgggat gcaaggctgg 19740
 ttcaacatac gcaaatcaat aaatgtaatc cagcatataa acagaaacaa agacaaaaac 19800
 cacatgatta tctcaataga tgcagaaaag gcatttgaca aaatttaaca actcttcacg 19860
 ctaaaaactc tcaatcaatt aggtattgat gggacgtatc tcaaaataat aagcactatc 19920
 tatgacaaac tcacagccaa tatcatactg aatgggcaaa aactggaagc attccctttg 19980
 aaaacgggca caagacaggg atgcctctc tcaccactcc tattcaacat agtggttgaa 20040
 gctctggcca gggcaattag gcaggagaag gaaataaagg gtattcaatt aggagaagag 20100
 gaagtcaaat tgtcctgtt tgcagatgac atgattgtat atctagaaaa ccccatcgtc 20160
 tcagcccaaa atctccttaa gctgataagc aacttcagca aagtctcagg atacaaaatc 20220
 aatgtacaaa aatcacaagc actcttatac atcaataaca gacaaacaga gagccaaatc 20280
 atgagtgaac tccattcac 20300

For purposes of the present invention, this DNA sequence will be referred to as SEQ ID NO:3. The location of the SNPs discussed further below is indicated by bold and larger font letters. Several additional sequences of DNA that are upstream from SEQ ID NO:3 are identified as relevant to the present invention. These DNA sequences are also found on NT_022030 and are

ggattaatca tgacaaaagt aatctaaatc tcgttaagac tacttaatga tcaatctttc 60
 cctctgtttt cctgactat agggaagtga attgccccaa tccttctcta tcacccccct 120
 gcagccatgc caatgcctta cctctgttat attcagccat aggggaagct tattctcata 180
 gaatcagggg ttggcatg**ta** gtcactagct attcttggtg agactagtga agatgagtga 240
 aggaaaatat tgcatagggtg aaatctcata ggcacaaata ggtgtttgtg agagtaacaa 300
 taaaagaaag tcattcccat actctagtag atgactcatt ttctcctcat tttttttttt 360

tcaaggcggt ctctacaacg gttaacctag taccaaaaat ccttctcttt tttcttgac 420

aaatcctggt caagtttagc tggcatttac tacgtccaag acattgtcca gatgctgtgg 480

For purposes of the present invention, this DNA sequence
will be referred to as SEQ ID NO:4.

agagaaagaa aggcaggcag caaggagaaa aaacattttt taaaaaaaga aaattaaaat 60

ccatgtaatg tctgatctct gttctgctgt atgtgtagat ctttccatat accaactcat 120

tagccttatt ttacagggtga ggaaaatgag acCgagagtc cttcttactt gaccaagttc 180

acacagcaag atcacacatg gtagaaccaa tgtagaacc taggtgtata cttgctcatt 240

caatatgtac aataattgca aaagtttcca taggtcttat tatatatcag gcactataaa 300

tgctatgcat gtgtcaacta atttaaacct aagcaatatt ataaggaagg tactattata 360

gaaatctcag ccttacaggt aagggaacag gaataaagag atgtgaggta atggcccaag 420

For purposes of the present invention, this DNA sequence
will be referred to as SEQ ID NO:5.

ataatctcct ttcaagtttt tatcctgtca cttgctagtt gtgtgatttg ggacaaatca 60

tttaactcct tgtaaaggga gagaagGaag gctgtaaaaa aattaagtaa taaaagata 120

aactccttgt ggtatatctt gttattgttc aaaaatattt attgcccttc ttaggatgtc 180

ttaggtcatt cttgcattgc tataaagaaa tacccaagtc tgggtaattt ataaagaata 240

gaggttaa at tggctcacag ttctgcaggc tgcacaggaa gcatcccact ggcgtctact 300

cacttctggt gaggactcag aaagcttttg cttatgacag caggctaagt gagagcaggt 360

For purposes of the present invention, this DNA sequence
will be referred to as SEQ ID NO:6.

[63] Several additional sequences of DNA that are downstream
from SEQ ID NO:3 are identified as relevant to the
present invention. These DNA sequences are also found
on NT_022030 and are

catggatatt ttactaccca ttgccttcta ggaaagggtg taacaaatag gaaatattaa 60

tatttttaaat gcctttgagg gtgttaaaaa gcacaactct aaggactggt tgtaaattc**C** 120
 aggtcaaagt ttgtttctcc ttctctatct cctaccttgg tgatggcctg atcttatatg 180
 gagtcactcc aactagaaac cacagaatca tccctagtgc ctacttctga ctcaactccat 240
 aactcaaaa gtcacctgac tctgcagaat ttctctagaa aaactctatg aaaacctatt 300
 cctgcctctc cacctgcata gatgtagctt catccaggct cttatgggtgc atggcctcgg 360
 ttactgcctt atcctttcta ctggcctctc aatctcccat ctgataccca ttaatgtact 420

For purposes of the present invention, this DNA sequence will be referred to as SEQ ID NO:7.

ccaaatactt ttaggcaca ctgggaagtt acattgtttc ttgcaagtga caggttgctc 60
 ttaattagt tctttctctc aaaaagagac tgctgactcc aaactgggaa gaaaccact 120
 caccagcaaa atgctgctga attcactctg atagtcttct aatctctcat cagtagatga 180
 caataatgaa gccagtattg ttaccacaag actcagatat **g**tctatcacc caagatgatt 240
 tctctttaag acgcaataaa agggaacttt tctcccccatt tattagcaac taagatgaaa 300
 tgagagccag agaaataaag tgaggaagga aagagaattt actaccttta caagctgaaa 360

For purposes of the present invention, this DNA sequence will be referred to as SEQ ID NO:8. In all upstream and downstream sequences (i.e. SEQ ID NOS: 4, 5, 6, 7, and 8), the location of SNPs are indicated by bold and larger font letters.

In situ hybridization

- [64] Double-stranded cDNA containing the RGS4 sequence was first amplified from normal human brain cDNA using custom designed primers (Forward primer sequence: CCGAAGCCACAGCTCCTC (SEQ ID NO: 3); Reverse primer sequence: CATCCCTCTCCCTTCAGGTG (SEQ ID NO: 4), and "touchdown" PCR with AmpliTaq Gold (PE Biosystems):

(94°C for 10 minutes (min), followed by 10 PCR cycles with a high annealing temperature 94°C for 30 seconds (sec), 62°C for 30 sec, and 72°C for 60 sec), 10 cycles with a medium annealing temperature (94°C for 30 sec, 60°C for 30 sec, 72°C for 60 sec), and 20 cycles at a low annealing temperature (94°C for 30 sec, 58°C for 30 sec, 72°C for 60 sec). The product of this touchdown PCR reaction produced a single bright band on a 2% agarose gel and was purified and ligated into a T/A plasmid cloning vector (AdvanTAge, Clontech) and transformed into competent *Escherichia coli* cells and plated overnight at 37°C. Colony PCR was performed on selected colonies containing the insert, and the products of these reactions were restriction digested and sequenced to verify orientation and insert identity.

[65] [³⁵S]-labeled riboprobes were synthesized using the T7 Riboprobe *In Vitro* Transcription System (Promega kit # P1460) and purified using RNeasy kit (Qiagen #74104). A scintillation counter was used to verify the specific radioactivity and yield of the probe. During hybridization, approximately 3 nanograms (ng) of probe was used per slide in a total volume of 90 µl. All other methods used were those described previously in Campbell et al., in *Exp. Neurol.* 160: 268-278, 1999, which is hereby incorporated by reference.

[66] Tissue blocks containing the regions of interest (PFC area 9, motor cortex [MC] and visual cortex [VC]) were identified using surface landmarks and sulci (the superior frontal gyrus, the central sulcus and precentral gyrus, and the calcarine sulcus, respectively). After histological verification of the regions, 20 µm sections containing these regions were

cut with a cryostat at -20°C , mounted onto gelatin-coated glass slides, and stored at -80°C until use. The slides were coded so that the investigator performing the analysis was blind to the diagnosis of the subjects.

[67] Following hybridization and washing, slides were air dried and exposed to BioMax MR film (Kodak) for 8-22 hours and then dipped in emulsion (NTB-2, Kodak), and exposed for 3-5 days at 4°C . High resolution scans of each film image were used for quantification of signal with Image (Scion Corporation, Fredrick, Maryland), version 4.0b), and darkfield images were captured from the developed slides. Throughout all steps and procedures, subject pairs were processed in parallel. Hybridization of sections with sense RGS4 riboprobe, used as a specificity control, did not result in detectable signal.

[68] Quantification was performed by subtracting the background white matter OD from the average signal OD measured in five non-overlapping rectangular regions on each section (3 sections per tissue block). In PFC and MC, these rectangular regions spanned cortical layers II-VI. Due to the lack of RGS4 signal in layer IV throughout the neocortex, and the great expansion of this layer in VC, the supragranular and infranular signal intensities were analyzed separately in VC. However, there were no significant differences in the levels of signal contained in the supra- and infragranular layers, so they were combined as a measure of overall VC signal intensity.

[69] Each *in situ* hybridization was repeated three times in separate hybridization reactions. The resulting ODs were background-corrected and averaged. Visual cortex

(V1) OD quantification, due to a bi-laminar transcript distribution, was performed separately for the supragranular and infragranular layers.

- [70] In order to search for novel candidate genes whose expression is consistently altered in schizophrenia, high-density cDNA microarrays (UniGEM-V, Incyte Genomics) were used to examine the expression patterns of over 7,800 genes and ESTs in post mortem samples of prefrontal cortex area 9 from six matched pairs of schizophrenic and control subjects.

Comparison and statistical analyses

- [71] As illustrated in FIG. 1B, a gene was determined to be expressed if the arrayed immobilized probe or target (the design of which is shown in FIG. 1A) was successfully amplified by PCR, produced a signal from at least 40% of the spot surface and had a signal/background ratio over 5-fold for either the cy3 or cy5 probe. Both images represent the same spot under cy3 and cy5 excitation, respectively. In this experiment, the balanced cy3 signal intensity (control or c-subject) was 6.2-fold brighter than the cy5 signal intensity (schizophrenic or s-subject).
- [72] Genes were comparably expressed between the control and experimental samples if the cy3/cy5 ratio or cy5/cy3 ratio was <1.6 . Over 80% of observations fell into this class. Gene expression was changed between the two samples at the 95% confidence level (95 % CL) if the cy3/cy5 or cy5/cy3 signal was 1.6 - 1.89. Gene expression was changed between the two samples at the 99% confidence level (99 % CL) if the cy3/cy5 or cy5/cy3 signal was 1.9.

[73] In the microarray analyses, data from experimental subjects were compared to data from matched control subjects in a pairwise design to control for the effects of age, race, sex and PMI on gene expression. To evaluate potential changes in gene group expression on the microarrays, two types of statistical measures were employed: 1) χ -square analysis was performed on the distribution of genes in a group versus the distribution of all genes called present on each individual microarray. The distribution of gene expression ratios was divided into five different bins based on confidence levels for individual gene comparisons: <-1.9 , -1.89 to -1.6 , -1.59 to 1.59 , 1.6 to 1.89 and >1.9 . 2) A paired t-test (degrees of freedom = 5) was used to compare mean expression ratios for a given gene group to the mean expression ratios for all expressed genes across all six subject pairs. A gene group was considered to be changed only if it reported differential expression by both the χ -square and t-test compared to the mean and distribution of all expressed genes. Microarray changes were also analyzed by descriptive statistics and correlation.

[74] To mimic the microarray comparisons, the *in situ* hybridization data were analyzed using ANCOVA with diagnosis as the main effect, subject pair as a blocking factor, and brain pH and tissue storage time as covariates. Furthermore, to verify that the pairing of subjects adequately controlled for sex, age, and PMI, we also conducted an ANCOVA with diagnosis as a main effect, and sex, age, PMI brain pH, and tissue storage time as covariates. Since both models produced similar results, the values from the ANCOVA with subject pair as a blocking factor are reported. Changes between groups

were also analyzed by descriptive statistics, Pearson correlation, and Factor analysis.

Pittsburgh cases and parents for genotyping analysis

- [75] Inpatients and outpatients were recruited at Western Psychiatric Institute and Clinic, a University of Pittsburgh-affiliated tertiary care center and 35 other treatment facilities within a 500 mile radius of Pittsburgh. The Diagnostic Interview for Genetic Studies (DIGS) was the primary source for clinical information for probands (Nurnberger, et al. *Archives of General Psych.* 51, 849-59; discussion 863-4, 1994). Additional information was obtained from available medical records and appropriate relatives, who also provided written informed consent. Consensus diagnoses were established by board certified psychiatrists. There were 93 Caucasian and 70 African-American cases. Genomic DNA, but not clinical information was available from all parents of the Caucasian cases. Cord blood samples were obtained from live births at Pittsburgh and served as unscreened, population-based controls. There were 169 individuals. They included 76 Caucasians and 93 African-Americans.

National Institute of Mental Health Collaborative Genetics Initiative (NIMH CGI) sample

- [76] From 1991-98, pedigrees having probands with schizophrenia or schizoaffective disorder, depressed (DSM IV criteria) were ascertained at Columbia University, Harvard University, and Washington University. The DIGS was the primary interview schedule. The families were ascertained if they included two or more affected first degree relatives (Cloninger

et al. *Am. J. Med. Gen.* **81**, 275-81, 1998, which is hereby incorporated by reference). We selected case-parent trios and available affected siblings from this cohort. Thus, 39 cases, their parents and 30 affected sibling-pairs were obtained. They comprised 25 Caucasian families, 10 who reported African-American ethnicity and 4 from other ethnic groups. Transmission disequilibrium test (TDT) analysis utilized only one case/family.

- [77] Written, informed consent was obtained from all participants. Ethnicity was based on self-report (maternal report for neonatal samples).

DNA sequencing and polymorphism detection

- [78] The genomic sequence for RGS4 was obtained from NT_022030 (390242 bp), a currently unfinished clone from Human Genome Project, Chromosome 1 database. The annotated data revealed three identified genes, namely, RGS4, MSTP032 and RGS5. The genomic organization of RGS4 and RGS5 includes 5 exons which is typical for the RGS family gene.
- [79] A panel of 10 African-American cases and 6 Caucasian controls was initially used to screen for polymorphisms in the exonic, intronic, and flanking genomic sequences of the RGS4 gene. The re-sequenced region included 6.8 kb upstream and 2.9 kb downstream of the coding sequence. The genomic sequence was used to design primers and amplicons ~500bp were generated, with overlapping sequences. The amplified fragments were sequenced using an ABI 3700 DNA sequencer. The sequencing panel that was used (n = 16) has over 80% power to detect SNPs with minor allele frequency over 5%

(Kruglyak et al. *Nature Gen.* **27**, 234-236, 2001, which is hereby incorporated by reference). We also sequenced cDNA sequences from the post-mortem samples reported on earlier (Mirnics et al. *Mol. Psychiatry* **6**, 293-301, 2001). The sequences were aligned using Sequencher (version 4.5) and polymorphisms were numbered consecutively. Additional SNPs localized to NT_022030 were obtained from the NCBI SNP database ("<http://www.ncbi.nlm.nih.gov/SNP>"). We also obtained genotype data from a prior study of the NIMH sample ("<http://zork.wustl.edu/nimh>").

Polymorphism analysis

- [80] PCR based assays included primers (5 pmol) with 200 μ M dNTP, 1.5 mM MgCl₂, 0.5 U of AmpliTaq Polymerase (PE Biosystems), 1x buffer and 60 ng of genomic DNA in 10 or 20 μ l reactions. The PCR conditions were 95°C for 10 min followed by 35 cycles (94°C for 45 sec, 60°C 45 sec and 72°C for 1 min). The final extension at 72°C for 7 min. The amplified products were digested with restriction endonucleases, electrophoresed on agarose gels, and visualized using ethidium stain. SNPs 4 and 18 were identified as single strand conformational polymorphisms (SSCP) (Orita et al. *DNAS* **86**, 2766-70, 1989). All genotypes were read independently by two investigators.
- [81] Polymorphisms were detected only in the intronic and flanking sequences of RGS4 (FIG. 6). Among 34 identified SNPs, one was selected from each of six sets which appeared to be in complete linkage disequilibrium in the re-sequenced panel. SNPs were further evaluated for informativeness (minor allele frequency > 0.1) and availability of reliable genotyping assays. Among the Caucasian cases from Pittsburgh, deviations from Hardy

Weinberg equilibrium (HWE) were noted for SNP 7 ($p < 0.03$) and SNP 13 ($p < 0.01$). Though all maternal genotypes conformed to HWE, deviations were noted at SNPs for the fathers of Pittsburgh cases at SNPs 4 and 18 ($p < 0.05$). For the analysis of IBD sharing among affected sibling-pairs from the NIMH samples, we also used genotypes for markers D1S1595, D1S484, D1S1677, D1S431 and D1S1589 (Faraone et al. *Am. J. of Med. Gen.* **81**, 290-5, 1998).

Statistical analysis

- [82] PEDCHECK software was used to check for Mendelian inconsistencies (O'Connell et al. *Am. J. of Hum. Gen.* **63**, 259-266, 1998, which is hereby incorporated by reference). χ^2 tests were employed for comparisons between cases and unrelated controls. We also used SNPEM software based on the EM algorithm to estimate and compare haplotype frequencies (Fallin, 2001, which is hereby incorporated by reference). We utilized GENEHUNTER software for TDT analysis of individual SNPs and haplotypes, as well as analysis of identity by descent among affected sibling-pairs (Kruglyak et al. *Am. J. of Hum. Gen.* **58**, 1347-63, 1996; Spielman et al. *Am. J. of Hum. Gen.* **54**, 559-60, 1994, both of which are hereby incorporated by reference). We also used TRANSMIT for global tests of association involving multiple haplotypes (Clayton et al. *Am. J. of Med. Gen.* **65**, 1161-1169, 1999a; Clayton et al. *Am. J. of Hum. Gen.* **65**, 1170-1177, 1999b, both of which are hereby incorporated by reference).

MICROARRAY RESULTS

- [83] Single gene transcripts were analyzed across all cDNA microarray comparisons. Across the six microarray comparisons over 90,000 data points were collected, and from these 44,000 were expression-positive observations, resulting in an average of 3,735 expressed genes/microarray. Of the expressed transcripts, 4.8% were judged to be differentially expressed (99% CL) between the schizophrenic and control subjects. The observed differences for any subject pair, in general, were comparably distributed in both directions: 2.6% of the genes were expressed at higher levels in schizophrenic subjects than in the matched controls, whereas 2.2% were expressed at lower levels in the schizophrenic subject.
- [84] Of all the expressed genes, RGS4 transcript reported the most significant decrease across all schizophrenic subjects. In fact, it was the only gene decreased at the 99% CL in all microarray comparisons. The microarray-bound, 571 base pair long, double-stranded cDNA immobilized probe corresponded to the 3' end of RGS4 and had a less than 50% sequence homology to any other known transcript, including RGS family members. This high binding specificity, coupled with strong cy3 and cy5 hybridization signal intensities, as shown in FIG. 1B, showed that RGS4 was robustly expressed in the human prefrontal cortex. Across the six microarray comparisons, RGS4 mRNA levels were decreased 50-84% in the PFC of schizophrenic subjects, as illustrated in FIG. 1C, while the expression of the ten other RGS family members represented on the microarray were unchanged in the schizophrenic subjects. In the scatter

plot shown in FIG. 1C, the X-axis reports subject pairs, the Y-axis reports percent change between schizophrenic and control subjects. Individual symbols represent a gene expression difference between a schizophrenic and control subject in a single pairwise comparison. The black dashed line denotes equal cy3 and cy5 signal intensity (similar expression) between schizophrenic and control subjects (0% change), green dashed line denotes the 95% confidence interval (37.5% change), red dashed line represents 99% confidence interval (47.5% change). Missing symbols in some pairwise comparisons indicate that the corresponding genes' microarray hybridization did not meet expression criteria. Across all the RGS members represented on the microarray, only RGS4 showed a consistent expression change over the 99% CL in schizophrenic subjects.

- [85] To confirm the microarray findings for the RGS4 expression changes, *in situ* hybridization was performed on the PFC from the same five subject pairs used for the microarray experiments (for pair 794c/665s, no sections were available from the same block of tissue used in the microarray experiment). As a further test of the robustness of the microarray data, five additional subject pairs were added to the *in situ* hybridization analysis. Radiolabeled cRNA probes designed against RGS4 mRNA were used to localize and quantify relative transcript levels. In the control subjects, RGS4 labeling was heavy in the prefrontal cortex, as shown in FIG. 2A, mimicking previously described labeling in the rat. In the gray matter of prefrontal cortex, the RGS4 riboprobe heavily labeled various size and shape cell profiles, including both projection neurons and interneurons. This labeling was the most prominent in

layers III and V, with sparse labeling in the intervening granular layer IV, and appeared to be present over both large pyramidal neurons and smaller cells that could represent interneurons. High power photomicrographs of PFC tissue sections from a schizophrenic (622s) and matched control (685c) subjects were viewed under darkfield illumination. Micrographs for each subject were taken under identical conditions. Roman numbers denote cortical layers. Pial surface is to the left. Strong labeling across all cortical layers except lamina IV was observed, and diminished labeling in the matched schizophrenic subject across all the layers was noted (scale bar = 400 μ m). White matter labeling was absent.

- [86] Based on optical density analysis, 9/10 subject pairs exhibited a 10.2% to 74.3% decrease in PFC RGS4 expression, as shown in FIG. 2B. The *in situ* hybridization data from 10 PFC pairwise comparisons were quantified using film densitometry. The X-axis represents subject classes, the Y-axis reports average film OD from 3 repeated hybridizations, measured across all layers. Lines connecting symbols indicate a matched subject pair. Note that in 10 PFC pairwise comparisons, 9 schizophrenic subjects showed RGS4 transcript reduction (mean = -34.5%; $F_{1,15} = 6.95$; $p = 0.019$).

Specificity of RGS4 expression changes

- [87] To investigate whether RGS4 transcript decrease is a specific alteration in schizophrenia, the same microarray data was analyzed for consistent gene expression changes across other RGS-family members (FIG. 1C). Nine of the eleven RGS family members represented with immobilized probes on the microarrays reported

expression in four or more microarray comparisons. RGS13, primarily lung-specific family member, was not expressed in any of the comparisons, while p115-RhoGEF reported expression in only one comparison. RGS4 was the only family member (and the only gene on the microarray) to report a consistent change in expression over the 99% CL in every schizophrenic subject. RGS5 mRNA (a gene also localized to cytogenetic position 1q21-22) was decreased at the 99%CL in one subject pair, at the 95% CL in another subject pair, and unchanged in the remaining 2 pairs that showed detectable RGS5 expression by microarrays. Expression of the other RGS family members did not display any consistent differences across the schizophrenic subjects. The mRNA from pair 567c/537s was analyzed a second time on the newest Incyte microarray, UniGEM-V2, which includes five additional RGS family members (RGSZ, RGS1, RGS7, RGS11, and RGS14). This analysis confirmed that, in the comparisons, RGS4 was the only significantly changed RGS family member.

- [88] Heterotrimeric G-proteins, the main substrates for RGS family members, were assessed for expression patterns. Several reports suggest $G\alpha$ changes associated with schizophrenia. Thus, it was desirable to assess whether the decrease in RGS4 expression correlated with changes in $G\alpha$ expression levels. Of the eight $G\alpha$ RGS substrates represented on the microarrays, only G_o expression was changed beyond the 95% CL in three or more pairwise comparisons. These three subjects with increased G_o levels (317s, 547s, and 622s) showed the most robust decrease in RGS4 expression both in the PFC microarray and *in situ* hybridization assays.

[89] Expression of 274 genes known to be involved in the G-protein signaling cascades (GPCR, heterotrimeric G-proteins, RGS, GIRKs, G-protein receptor kinases, and mitogen-activated protein kinases) were analyzed in a gene group comparison. An average of 105 genes belonging to this group were expressed in each comparison. The results of microarray analyses showing G-protein and 1q21-22 locus-related expression differences in the PFC of six pairs of schizophrenic and control subjects are shown in FIGS. 3A and 3B. For both gene groups, all expressed genes were classified into signal intensity difference intervals (0.1 bins) according to their cy5/cy3 signal ratio. Transcripts in a "1" bin had identical cy5 vs. cy3 signal intensities. Positive values (to the right) on the X-axis denote higher cy5 signal in schizophrenic subjects ($S > C$), negative values (to the left) correspond to higher cy3 signal intensity in the control subjects ($C > S$). The Y-axis reports percentage of expressed genes across the six subject pairs per bin for each gene group. In both panels, the white bars (*All genes*) denote distribution of all expressed genes across the six PFC pairwise comparisons ($n = 22,408$). Additionally, in both panels, RGS4 contribution to the transcript distribution is denoted by a hatched bar. Note that in both FIG. 3A and FIG. 3B, the cy3/cy5 signal distribution of G-protein and 1q21-22 gene groups was comparable to the distribution of all expressed genes across the six microarray comparisons.

[90] At the 99% confidence level, 5.6% of G-proteins showed a different distribution between schizophrenic and control subjects, as shown in FIG. 3A: 2.8% of G-proteins were decreased, while 2.8% were increased in the PFC of

schizophrenic subjects. Of the 2.8% decrease in schizophrenic subjects, RGS4 observations alone accounted for nearly half of the decrease. When RGS4 was removed from the G-protein group, a gene group analysis by χ^2 test and t-test closely matched the distribution of all expressed genes, suggesting that the majority of different expression levels can be attributed to normal human variability. Except RGS4, no other member of the G-protein gene group was consistently changed across the subject pairs over the 95% or 99% confidence levels.

- [91] The RGS4 gene has been mapped to locus 1q21-22, a novel schizophrenia locus recently implicated by pedigree studies with a linkage of disease score (LOD) of 6.5 as described by Brzustowicz et al. *supra*. To address if any other genes at this locus displayed altered expression in the PFC of schizophrenic subjects, 70 additional transcripts originating from this cytogenetic region were analyzed. At the 99% CL, 0.4% of 1q21-22 genes were increased, and 5.9% were decreased in the schizophrenic subjects. Of the transcripts decreased in schizophrenic subjects, RGS4 observations alone accounted for nearly half of the decreases, as shown in FIG. 3B. Furthermore, of all the genes on the 1q21-22 locus, only RGS4 showed a consistent expression change across all the pairwise comparisons over the 95% or 99% confidence levels. Of the remaining genes on this locus, only the ALL1-FUSED gene (AF1q GenBank Accession #U16954) reported consistent expression change over the 95% CL in the schizophrenic subjects in three or more pairwise comparisons. Furthermore, as a gene group, the expression of the remaining genes on locus 1q21-22 showed the same overall pattern as genes located on non-

schizophrenia loci or the overall average gene expression which is shown in FIG. 3B.

Regional RGS4 gene expression changes

- [92] To test whether RGS4 transcript decrease is specific to the prefrontal cortex or includes a more widespread cortical deficiency, RGS4 expression was assessed by *in situ* hybridization in the visual cortex (VC) and motor cortex (MC) from the same 10 pairs of control and schizophrenic subjects (for pair 558c/317s MC material was not available, and this pair was substituted with pair 794c/665s). The figure layout for FIG. 4A-D is similar to that of FIG. 2A-B. In VC, RGS4 *in situ* hybridization showed heavy labeling under darkfield illumination of diverse cell population in the gray matter, with a very prominent bi-laminar labeling pattern in the supragranular and infragranular layers, as shown in FIG. 4A. Roman numbers denote cortical layers, scale bar = 400 μ m. There was very sparse labeling in the well-developed layer IV, with very few cellular elements exhibiting detectable levels of RGS4 mRNA. These high power photomicrographs show that RGS4 levels are significantly decreased in the VC region of the schizophrenic subjects. The OD measurements on these two layers were performed separately.
- [93] Across the same ten pairwise comparisons that were examined in the PFC hybridizations, combined RGS4 expression in supragranular and infragranular layers of VC was decreased by 32.8% ($F_{1,15} = 8.24$; $p = 0.012$) as shown in FIG. 4B.
- [94] In MC, RGS4 expression was concentrated over the cell-rich layers I-III and V-VI of both control and

schizophrenic subjects, as shown in FIG. 4C. High power photomicrographs of MC tissue sections from the same matched pair of schizophrenic and control subject are represented in FIG. 2A and FIG. 4A, viewed under darkfield illumination. Roman numbers denote cortical layers, scale bar = 400 μ m. Because of the attenuated layer IV in motor cortex, the RGS4 labeling is almost uniform across all layers.

[95] Similar to the RGS4 transcript decrease observed in supragranular VC, schizophrenic subjects across the same 10 subject pairs were analyzed in MC. The mean RGS4 expression in MC shown in FIG. 4D, measured across all the layers, was decreased by 34.2% across the 10 schizophrenic subjects ($F_{1,15} = 10.18$; $p = 0.006$).

[96] In the PFC, VC, and MC of subjects with schizophrenia, RGS4 expression was consistently decreased compared to the PFC of subjects with the diagnosis of MDD, as shown in the schematic of FIG. 5. In contrast, factor analysis of the pairwise differences in RGS4 gene expression across 3 different cortical areas for all 9 common schizophrenic and control subject pairs revealed that over 84% of the total variance in expression was accounted for by diagnosis (variance proportion = 0.848, eigenvalue = 2.544, $p = 0.001$). The X-axis represents experimental groups, the Y-axis reports percent RGS4 expression change in PFC, VC, MC, in schizophrenic subjects (SCH) and PFC of subjects with MDD viewed by *in situ* hybridization. Each symbol represents percent of change between a single pairwise comparison; same symbols represent the same subject pairs. Arrows represent mean expression difference for each group. The same schizophrenic subjects showed a comparable and

highly correlated decrease in RGS4 expression across all three cortical regions (PFC-VC: $r = 0.88$, $p = 0.0003$; PFC-MC: $r = 0.69$, $p = 0.0384$; VC-MC: $r = 0.76$, $p = 0.0144$). In contrast, subjects with MDD reported variable RGS4 expression changes when compared to their matched controls.

[97] The combined data indicate that RGS4 transcript changes are a result of the pathophysiological changes related to schizophrenia and not due to confounds. Furthermore, the RGS4 expression decrease appears to be specific and unique to schizophrenia, and not a hallmark of the major depressive disorder.

[98] RGS4 labeling in the white matter was comparable to background labeling across all brain regions, suggesting that RGS4 is primarily expressed in neuronal cells. The labeling was abundant in the majority of interneurons and projection neurons. However, in some pyramidal cells and interneurons RGS4 labeling could not be detected. RGS4 labeling was heavy in all cortical layers, except layer IV, where RGS4 expression was both sparse and light. This overall pattern of labeling was comparable across all three cortical regions (PFC, VC, MC). As the granular layer IV is the widest in the primary visual cortex, in this region RGS4 labeling was prominent in supragranular and infragranular layers, separated by a wide zone of mostly unlabeled granular cells. The overall distribution pattern of the RGS4 message does not mimic the known expression patterns of neurotransmitter systems, suggesting that RGS4 regulates many functionally distinct neuronal populations.

[99] Together, the microarray and *in situ* hybridization methods suggest decreased RGS4 expression is a

consistent characteristic of schizophrenic subjects. Several causes of the reduced RGS4 expression may be offered, including adaptive and genetic changes in schizophrenic patients. It was hypothesized that reduction in RGS4 expression was generated by alterations in the RGS4 gene. In addition, it was contemplated that variations in the DNA upstream and downstream from the coding region of the RGS4 gene may also impact the expression of the RGS4 transcript. These possibilities were investigated by searching for SNPs in the RGS4 gene.

- [100] The specificity of the reduced expression of RGS4 message for schizophrenic patients was confirmed in a series of control experiments. The same reduced level of RGS4 message was not observed in patients suffering from major depressive disorder. In addition, prolonged treatment of non-human primates with the anti-psychotic haloperidol did not result in decreased levels of RGS message in the cerebral cortex. This result indicates that chronic exposure to anti-psychotic drugs are unlikely to be responsible for the depressed levels of RGS4 message observed in schizophrenic patients.

GENOTYPING RESULTS

- [101] 34 single nucleotide polymorphisms (SNPs) were identified after re-sequencing all exons, introns and flanking 5' and 3' UTRs of the RGS4 coding region (FIG. 6). Thirteen SNPs were chosen for analysis using the TDT. SNPs are explicitly defined in Table 1. When the SNPs were tested individually, significantly increased transmission at SNP4 was observed in the Pittsburgh sample. 'Moving window' haplotype analyses using two to four contiguous SNPs, revealed significant association

for several haplotypes; all but one included SNPs 1, 4, 7, or 18 (Table 2). A global test of association for haplotypes encompassing these SNPs was significant (TRANSMIT software, $\chi^2 = 16.6$, 8 df, $p = 0.035$). There were 39 cases with schizoaffective disorder in the sample; these trends remained significant when the sample was restricted to individuals with schizophrenia ($\chi^2 = 13.0$, 6 df, $p = 0.043$).

[102] TDT analysis was conducted next in the ethnically diverse NIMH sample using the same set of SNPs. Significant transmission distortion was observed individually at SNPs 1, 4 and 18 (Table 2). Exclusion of African-American families from the sample also

SNP #	Location of the SNP within the SEQ	Nucleotide identity in SEQ ID NO:3	Observed Nucleotide variation
27,859	199 {SEQ ID NO:4}	T	C
34,653	153 {SEQ ID NO:5}	C	T
90,387	87 {SEQ ID NO:6}	G	A
SNP1	4121 {SEQ ID NO:3}	C	T
SNP2	4123 {SEQ ID NO:3}	T	A
SNP3	4368 {SEQ ID NO:3}	A	C
SNP4	4621 {SEQ ID NO:3}	A	C
SNP5	4790 {SEQ ID NO:3}	C	T
SNP6	4816 {SEQ ID NO:3}	G	T
SNP7	4970 {SEQ ID NO:3}	C	T
SNP8	5055 {SEQ ID NO:3}	C	G
SNP9	5295 {SEQ ID NO:3}	G	A
SNP10	5695 {SEQ ID NO:3}	G	A
SNP11	7375 {SEQ ID NO:3}	G	T
SNP12	7759 {SEQ ID NO:3}	G	A
SNP13	8596 {SEQ ID NO:3}	G	A
SNP14	9603-9609 {SEQ ID NO:3}	AGTTTGG	7 bases Absent
SNP15	9892 {SEQ ID NO:3}	C	A
SNP16	9963 {SEQ ID NO:3}	C	A
SNP17	10132 {SEQ ID NO:3}	G	A
SNP18	11056 {SEQ ID NO:3}	T	C
SNP19	11091 {SEQ ID NO:3}	C	T
SNP20	11106 {SEQ ID NO:3}	C	A
SNP21	11774 {SEQ ID NO:3}	G	T
SNP22	12143 {SEQ ID NO:3}	G	A
SNP23	12145 {SEQ ID NO:3}	G	T
SNP24	14367 {SEQ ID NO:3}	A	G
SNP25	17028 {SEQ ID NO:3}	A	Base absent
SNP26	17630 {SEQ ID NO:3}	G	T
118740	120 {SEQ ID NO:7}	C	G
130121	221 {SEQ ID NO:8}	G	C

[103] **Table 1.** Location of single nucleotide polymorphisms relevant to the present invention. The location of the SNP within the sequence is listed as is the variation observed in the collected samples. SNP 14 is the absence of the listed 7 bases at the indicated location.

revealed significant results for these SNPs ($p = 0.023$, 0.011 and 0.033 respectively). However, the transmitted alleles differed from the Pittsburgh sample. Moving window haplotype analyses revealed preferential transmission for more extensive chromosomal segments than the Pittsburgh sample. Like the Pittsburgh sample, all but one of haplotypes with significant increased transmission included SNPs 1, 4, 7 or 18. A global test for association was also significant for haplotypes encompassing these SNPs (TRANSMIT analysis; $\chi^2 = 18.8$, $p = 0.016$, 8 df).

[104] If the significant TDT results were due to linkage, it was reasoned that the affected sibships in the NIMH sample should yield evidence for increased haplotype sharing. For 30 available affected sib-pairs, the proportion of 0, 1, or 2 haplotypes identical by descent (IBD) were elevated over expectations of 0.25, 0.50, 0.25; namely 0.11, 0.44, 0.45 respectively (for SNPs 1, 4, 7 and 18 analyzed in conjunction with 5 flanking short tandem repeat polymorphisms genotyped previously). Increased IBD sharing was also observed when these sets of SNPs or STRPs were analyzed separately.

[105] Association at the population level was assessed by comparing Caucasian cases from each sample separately with two independent groups of Caucasian community-based controls. Since SNPs 1, 4, 7 and 18 appeared to be critical for transmission distortion in both samples, genotypes and allele frequencies for these SNPs were analyzed. Haplotypes frequencies were estimated using an expectation-maximization algorithm (EM), paying particular attention to haplotypes VI and XI, the haplotypes with excess transmission in the NIMH and

Pittsburgh samples, respectively (Table 3). SNP 14 was informative only among African-Americans and so was analyzed separately using 70 African-American cases and 93 control individuals from Pittsburgh. Significant case-control differences were not noted for any of the comparisons. The failure to detect association may reflect superior power for the TDT in the context of population sub-structure.

No.	Haplotype	Neonatal Controls	Adult controls	Pittsburgh Cases	NIMH Cases
	SNP 1—4—7—18				
I	o—o—o—o	0.096	0.066	0.078	0.067
II	•—o—o—o	0.004	0.021	0.022	0.083
III	o—•—o—o	0.006	0.006	0.000	0.000
IV	•—•—o—o	0.000	0.000	0.000	0.000
V	o—o—•—o	0.000	0.000	0.006	0.000
VI	•—o—•—o	0.388	0.442	0.378	0.392
VII	o—•—•—o	0.000	0.006	0.000	0.000
VIII	•—•—•—o	0.000	0.000	0.006	0.000
IX	o—o—o—•	0.000	0.004	0.000	0.017
X	•—o—o—•	0.000	0.006	0.000	0.000
XI	o—•—o—•	0.439	0.425	0.494	0.417
XII	•—•—o—•	0.008	0.013	0.000	0.000
XIII	o—o—•—•	0.000	0.000	0.000	0.000
XIV	•—o—•—•	0.053	0.013	0.017	0.025
XV	o—•—•—•	0.006	0.000	0.000	0.000
XVI	•—•—•—•	0.000	0.000	0.000	0.000

[106] **Table 2.** Haplotype based comparisons among cases and unrelated controls. The Caucasian cases from Pittsburgh (n = 93) and NIMH (n = 25) were compared separately with unscreened Caucasian controls from Pittsburgh (n = 76). Bonferoni corrections have been applied for the Pittsburgh case-control comparisons, but not for comparisons involving the NIMH cases. An omnibus test based on likelihood ratios was used to estimate overall differences in haplotype frequencies (Fallin et al., Gen. Res. 11, 143-51, 2001) and was significant for both comparisons ($\chi^2 = 88.7$, $p < 0.0001$ and $\chi^2 = 30.1$, $p < 0.0003$ respectively for Pittsburgh and NIMH cases). Similar significant differences based on 3 SNP haplotypes were present, but are not shown. For each SNP, 'o' represents allele 1 and '•' represents allele 2. OR - Odds ratio; NS - Not significant.

SNP	27859	90387	snp1	snp4	snp7	snp18	snp23	118740	130121
27859		0.096	0.064	0.076	0.287	0.009	0.000	0.000	0.000
90387	0.132		0.000	0.000	0.000	0.000	0.000	0.001	0.627
snp1	-0.123	-0.501		0.000	0.000	0.000	0.000	0.450	0.477
snp4	0.101	-0.501	-1.000		0.000	0.000	0.000	0.128	0.515
snp7	-0.075	0.783	0.970	-0.961		0.000	0.000	0.012	0.068
snp18	0.177	0.377	-0.677	0.989	-0.961		0.000	0.000	0.041
snp23	0.527	-0.302	-1.000	1.000	-0.847	0.674		0.499	0.002
118740	0.385	0.163	0.048	-0.083	0.172	-0.233	0.042		0.000
130121	-0.505	0.049	-0.059	0.046	-0.163	0.174	-0.154	-0.956	

[107] **Table 3.** Pair-wise linkage disequilibrium between SNPs at RGS4. Population based control individuals (n = 76) were used to estimate linkage disequilibrium. The figures above the diagonal represent D' and estimates for statistical significance (p values) are below the diagonal.

[illegible]

[108] **Table 4.** SNPs and Haplotypes at RGS4 with increased transmission distortion. TDT analysis of case-parent trios included 93 families from Pittsburgh and 39 families from the NIMH cohort. Only statistically significant increased transmissions are shown. The shaded haplotypes correspond to haplotypes VII and X, respectively from Table 2. T/NT-Transmitted/not transmitted; o-Allele 1, •-Allele 2 at each SNP;/-Allele not specified at this locus; *p<0.05, **p<0.01, *** p<0.005.

[109] The demonstration of the association between these SNPs and schizophrenia offers a large number of applications in the diagnostic and therapeutic fields. Thus, embodiments of the present invention offer the possibility of diagnosing schizophrenia by means of a biological test and no longer exclusively by means of clinical evaluations. Embodiments of the present invention can also be applied to diagnosing pathologies of the schizophrenia spectrum, such as, in particular, schizotypy, schizoid individuals, etc. Embodiments of the present invention make it possible to refine the criteria for diagnosing these pathologies, which is currently entirely established clinically. Furthermore, embodiments of the invention also makes it possible to demonstrate susceptibility to schizophrenia by means of identifying a genetic vulnerability in the families of patients who posses the identified SNPs in the RGS4 coding region and flanking regions. Once individuals have been identified as being susceptible to schizophrenia, the utility of prophylactic treatment may be investigated.

[110] The DNA sample to be tested can be obtained from cells that have been withdrawn from the patient. These cells are preferably blood cells (e.g. mononucleated cells), that are easily obtained by the simple withdrawal of blood from the patient. Other cell types, such as fibroblasts, epithelial cells, keratinocytes, etc., may also be employed. The DNA may then extracted from the cells and used to detect the presence of SNPs in the RGS4 coding region and flanking regions.

[111] Most preferably, the DNA extract is initially subjected to one or more amplification reactions in order to

obtain a substantial quantity of material corresponding to the region carrying the RGS4 coding region and flanking regions. The amplification can be achieved by any technique known to the skilled person, and in particular by means of the so-called PCR technique as described above. To this end, embodiments of the present invention also relate to specific primers which make it possible to amplify DNA fragments that are of small size and which carry the RGS4 gene, flanking regions thereof, or portions thereof generated from SEQ ID NOS. 3, 4, 5, 6, 7, or 8. Portion of a polynucleotide sequence is specifically intended to refer to any section of SEQ ID NOS. 3, 4, 5, 6, 7, or 8 that can be used in the practice of this invention, such as use as a primer to identify the presence of SEQ ID NOS. 3, 4, 5, 6, 7, or 8 or variations thereof in a patient or a section of SEQ ID NOS. 3, 4, 5, 6, 7, or 8 that can be used to amplify the entire sequence. The phrase contiguous portion is meant to refer to a series of bases that are adjacent to one another within a polynucleotide sequence. In the context of the present invention, the word gene is intended to mean the protein coding region, the proximal 5' and 3' untranslated regions, as well as any distal and proximal regulatory domains. The phrase gene-coding region is meant to refer to the stretch of DNA that begins at the transcription initiation site and includes all exonic and intronic sequences that encode a protein.

- [112] Embodiments of the present invention may also involve isolating DNA sequences and ligating the isolated sequence into a replicative cloning vector which comprises the isolated DNA of the RGS4 gene, based upon or derived from the cDNA of SEQ ID NOS. 3, 4, 5, 6, 7,

or 8 and a replicon operative in a host cell. Additional embodiments include an expression system which comprises isolating DNA of the RGS4 gene, based upon complementarity to SEQ ID NOS. 3, 4, 5, 6, 7, or 8 and operably linking this DNA to suitable control sequences. Recombinant host cells can be transformed with any of these replicative cloning vectors and may be used to overproduce the RGS4 protein.

[113] Embodiments of the present invention also include kits that will facilitate the diagnosis of schizophrenia through the amplification of segments of the 1q21-22 locus. Several methods providing for this amplification are described including: at least a pair of single-stranded DNA primers wherein use of said primers in a polymerase chain reaction results in amplification of a portion of the RGS4 gene fragment, wherein the sequence of said primers is derived from the regions of the cDNA defined by or complementary to SEQ ID NOS: 1, 3, 4, 5, 6, 7, or 8. Similarly, embodiments of the invention also provide for a pair of single-stranded DNA primers wherein use of said primers in a polymerase chain reaction results in amplification of an RGS4 gene fragment, wherein the sequence of said primers is based on the exon regions of chromosomal DNA derived from SEQ ID NOS:1 or 3.

[114] Various nucleic acid probes and primers specific for RGS4 (derived from or complementary to SEQ ID NOS. 3, 4, 5, 6, 7, or 8) may also be useful in diagnostic and therapeutic techniques and are included within the present invention. Among these are a nucleic acid probe complementary to portions or the entirety of human RGS4 gene as well as a nucleic acid probe complementary to

human altered RGS4 gene sequences wherein said nucleic acid probe hybridizes to a variant of the RGS4 gene under hybridization conditions which prevent hybridizing of said nucleic acid probe to a wild-type RGS4 gene. Probes that are complementary to portions or the entirety of the RGS4 coding region and flanking regions that contain SNPs may also be used in these diagnostic tests. Any primer which makes it possible to amplify a fragment of the RGS4 coding region or flanking regions also forms part of the present invention. The primers that are used within the context of the invention can be synthesized by any technique known to the skilled person. The primers can also be labeled by any technique known to the skilled person.

- [115] The invention may also be practiced through detection of SNPs in the RGS4 coding region or flanking regions by a variety of techniques. The techniques which may preferably be employed are DNA sequencing and gel separation.
- [116] Any sequencing method known to the skilled person may be employed. In particular, it is advantageous to use an automated DNA sequencer. The sequencing is preferably carried out on double-stranded templates by means of the chain-termination method using fluorescent primers. An appropriate kit for this purpose is the Taq Dye Primer sequencing kit from Applied Biosystem (Applied Biosystem, Foster City, CA). Sequencing the SNPs in the RGS4 coding region and the flanking regions makes it possible to identify directly the SNPs that are present in the patient.
- [117] An additional preferred technique for demonstrating the SNPs in the RGS4 coding region and flanking regions is

that of separation on a gel. This technique is based on the migration, under denaturing conditions, of the denatured DNA fragments in a polyacrylamide gel. The bands of DNA can be visualized by any technique known to the skilled person, with the technique being based, such as by using labeled probes that are complementary to the entirety or portions of the RGS4 coding region and flanking regions. Alternatively, the bands may be visualized by using ethidium bromide or else by means of hybridization with a radiolabeled probe.

[118] In addition, measuring the expression of RGS4 message in peripheral tissue allows the diagnosis and determination of the susceptibility to schizophrenia in humans. As a matter of convenience, the reagents employed in the present invention can be provided in a kit packaged in combination with predetermined amounts of reagents for use in determining and/or quantifying the level of RGS4 expression. For example, a kit can comprise in packaged combination with other reagents any or all of the following components: appropriate detectors, buffers, deoxynucleotide triphosphates, ions provided by $MgCl_2$ or $MnCl_2$, and polymerase(s). The diagnostic kits of the invention may further comprise a positive control and/or a negative control as well as instructions for quantitating RGS4 expression.

[119] Additionally, an embodiment of the present invention relates to ascertaining levels of the RGS4 protein. The level of RGS4 protein can be detected by analyzing binding of a sample from a subject with an antibody capable of binding to RGS4. An embodiment of this detection method utilizes an immunoassay. The sample from a subject may preferably be a biopsy of skeletal

muscle, though any tissue accessible to biopsy may be used.

[120] In addition to providing generally useful diagnostic kits and methods, embodiments of the present invention may provide a method for augmenting traditional treatments by supplying the RGS4 protein to a subject and/or augmenting the subject's medication, such as antipsychotic drugs, and providing an improved therapeutic outcome.

[121] Further embodiments of the present invention may relate to the construction of an animal model of schizophrenia. Transgenic mice technology involves the introduction of new or altered genetic material into the mouse germ line by microinjection, retroviral infection or embryonic stem cell transfer. This results in lineages that carry the new integrated genetic material. Insertional mutagenesis occurs when integration of the microinjected genetic material into the host genome alters an endogenous gene resulting in a mutation. Methods of transferring genes into the germline, the expression of natural and hybrid genes and phenotypic changes that have occurred in transgenic mice are described by Palmiter and Brinster in Ann. Rev. Genet. 20 (1986) 465-499. Methods of foreign gene insertion, applications to foreign gene expression, and the use of transgenic mice to study immunological processes, neoplastic disease and other proliferative disorders are described by Gordon in Intl. Rev. Cytol. 115, 1989, 171-299 both of which are hereby incorporated by reference. A further example of genetic 'knock-in' technology may be found in Nebert, et al., Ann. N.Y. Acad. Sci. 919, 2000, 148-170 which is hereby incorporated by reference. The insertion of SEQ

ID NO:3 containing some or all of the described SNPs into a mouse germ line may be expected to result in adult mice that may be used as an experimental model of schizophrenia. The insertion of SEQ ID NO:3 containing one or more of the variations listed in Table 1 with standard on:off regulatory domains will allow for the creation of mice deficient in RGS4 expression at specified times, and may be used as an experimental model of schizophrenia.

[122] Having now fully described embodiments of the present invention, it will be appreciated by those skilled in the art that the same can be performed within a wide range of equivalent parameters, concentrations, and conditions without departing from the spirit and scope of the invention and without undue experimentation. While this invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This application is intended to cover any variations, uses, or adaptations of the invention.